

Fan, Montemurro, Motakis, Praveen, Rusonik, Skoufranis and Tobin

*Restricted invertibility of continuous matrix
functions*

OPERATORS AND MATRICES

AIMS AND SCOPE

'Operators and Matrices' (OaM), aims towards developing a high standard international journal which will publish top quality research papers in matrix and operator theory and their applications. The journal will publish mainly pure mathematics, but occasionally papers of a more applied nature could be accepted providing they have a nontrivial mathematical content. OaM will also publish expository papers and relevant book reviews.

OaM is published quarterly, in March, June, September and December.

SUBMISSION

Manuscripts should be submitted through OaM page: oam.ele-math.com, by e-mail of Editorial office, or directly to any one of the editors working in a related subject area.

Manuscripts are accepted for refereeing on the understanding that the same work has not been published (except in the form of an abstract), and is not under consideration for publication elsewhere.

Authors are advised to send notification of submission by e-mail to oam@element.hr (author(s), title, number of pages, name of editor to whom paper was sent to).

In order to facilitate refereeing, copies of those papers (whether by the author or someone else) which are essential and referred to in the manuscript but are not conveniently accessible, should be enclosed.

The publisher strongly encourages submission of manuscripts written in TeX or one of its variants LaTeX, AMSTeX or AMSLaTeX. On acceptance of the paper, authors will be asked to send the file by electronic mail. The file must be the one from which the accompanying manuscript (finalized version) was printed out. A poscript or pdf file of this final version must also be enclosed.

TITLE PAGE

The following data should be included on title page: the title of the article, author's name (no degrees) author's affiliation, e-mail addresses, mailing address of the corresponding author, and running head less than 60 characters.

ABSTRACT, KEY WORDS, SUBJECT CLASSIFICATION

The manuscript must be accompanied by a brief abstract, no longer than 100-150 words. It should make minimal use of mathematical symbols and displayed formulas. Mathematics Subject Classification (2010) and a list of 4-5 key words must be given.

FIGURES

Figures should be prepared in a digital form suitable for direct reproduction, at resolution of 300 dpi or higher, and in EPS, TIFF or JPEG format.

REFERENCES

Bibliographic references should be listed alphabetically at the end of the article. The authors should consult Mathematical Reviews for the standard abbreviations of journal names.

PROOFS, PAGE CHARGES, OFFPRINTS

The galley proofs will be sent to corresponding author. Late return of the proofs will delay the article to a later issue.

There are no page charges. Authors will receive PDF file of the printed article free of charge. Additional offprints may be ordered from OaM prior to publication.

FORTHCOMING PAPERS

Papers accepted and prepared for publication will appear in the forthcoming section of Journal Web page. They are identical in form as final printed papers, except volume, issue and page numbers.

OaM is published by Publishing House **ELEMENT**, Zagreb, Croatia.

The publication of this journal is supported by Ministry of science, education and sports, Croatia

All correspondence and subscription orders should be addressed to the Editorial Office:

www.ele-math.com

e-mail: oam@ele-math.com

Operators and Matrices
Editorial Office
Menceticeva 2, 10000 Zagreb, Croatia
Fax: +385 1 6008799

RESTRICTED INVERTIBILITY OF CONTINUOUS MATRIX FUNCTIONS

ADRIAN FAN, JACK MONTEMURRO, PAVLOS MOTAKIS, NAINA PRAVEEN,
ALYSSA RUSONIK, PAUL SKOUFRANIS AND NOAM TOBIN

(Communicated by I. Klep)

Abstract. Motivated by an influential result of Bourgain and Tzafriri, we consider continuous matrix functions $A : \mathbb{R} \rightarrow M_{n \times n}$ and lower ℓ_2 -norm bounds associated with their restriction to certain subspaces. We prove that for any such A with unit-length columns, there exists a continuous choice of subspaces $t \mapsto U(t) \subset \mathbb{R}^n$ such that for $v \in U(t)$, $\|A(t)v\| \geq c\|v\|$ where c is some universal constant. We provide two methods. The first relies on an orthogonality argument and it yields an optimal asymptotic dependence for $\dim(U(t))$ on n and $\sup_{t \in \mathbb{R}} \|A(t)\|$ but it does not preserve any structure for $U(t)$. The second is probabilistic and combinatorial in nature and it does not yield the optimal bound for $\dim(U(t))$ but the $U(t)$ obtained in this way are guaranteed to have a canonical representation as joined-together spaces spanned by subsets of the unit vector basis.

1. Introduction

This paper concerns itself with continuous matrix functions $A : \mathbb{R} \rightarrow M_{n \times n}$ restricted to linear subspaces and certain lower ℓ_2 -norm bounds satisfied on these subspaces. It is motivated by specific results in the non-continuous (static) setting. Inspired by problems in harmonic analysis and the geometry of Banach spaces, a seminal 1987 article by Bourgain and Tzafriri ([5]) proved the following.

THEOREM 1.1. ([5]) *There exist universal constants $c_0, d_0 > 0$ such that for any $A \in M_{n \times n}(\mathbb{R})$ with $\|Ae_i\| = 1$ for $1 \leq i \leq n$ there exists $\sigma \subset \{1, \dots, n\}$ with $|\sigma| \geq d_0 n \|A\|^{-2}$ such that for any set of scalars $\{a_j\}_{j \in \sigma}$,*

$$\left\| \sum_{j \in \sigma} a_j A e_j \right\|_2 \geq c_0 \left(\sum_{j \in \sigma} |a_j|^2 \right)^{1/2}. \quad (1)$$

Equivalently, for $U_\sigma = \langle e_j : j \in \sigma \rangle$ and for any $v \in U_\sigma$, $\|Av\| \geq c_0 \|v\|$.

Mathematics subject classification (2020): 46B07, 46B20, 47A68, 15A09.

Keywords and phrases: Restricted invertibility, continuous matrix functions.

This research was conducted during a Fields Undergraduate Summer Research Program in 2021 and the first, second, fourth, and fifth authors were supported by this program. The third author was supported by NSERC Grant RGPIN-2021-03639. The sixth author was supported by NSERC Grant RGPIN-2017-05711. The seventh author was supported by an Undergraduate Student Research Award (NSERC).

For the remainder of the paper, let c_0 and d_0 be a pair of constants satisfying the conclusion of Theorem 1.1. We point out that a dimensional estimate of order $n\|A\|^{-2}$ is optimal up to a constant (see Remark 3.7). Intuitively, the Bourgain-Tzafriri result guarantees the existence of a “large” coordinate subspace of \mathbb{R}^n on which A does not shrink vectors “excessively.” The above result implies that A is invertible when restricted to U which was the original motivation for [5], and hence the term restricted invertibility. The U from the above theorem is a particularly simple type of subspace namely one spanned by a subset of standard basis vectors.

The coordinate structure of the subspace U is a central aspect of Theorem 1.1. A far more elementary spectral decomposition argument also yields a high-dimensional subspace U , without any structure, on which A satisfies a lower ℓ_2 bound. For any $A \in M_{n \times n}(\mathbb{R})$ and any $\gamma \in (0, 1)$ there exist a subspace U of \mathbb{R}^n with such that for any $x \in U$, $\|Ax\| \geq \gamma\|x\|$ and

$$\dim(U) > \left(\frac{\|A\|_{\text{HS}}^2}{n} - \gamma^2 \right) n\|A\|^{-2}. \quad (2)$$

If A has unit-length columns then $\|A\|_{\text{HS}}^2 = n$ and thus $\dim(U) > (1 - \gamma^2)n\|A\|^{-2}$ (see Section 2).

Theorem 1.1 from [5] and subsequent work of Bourgain and Tzafriri ([6], [7]) are strongly related to the famous Kadison-Singer conjecture [10]. This was a central problem in C^* -algebras that was restated by Anderson in [2] as a problem about matrices and it was proved by Casazza and Tremain in [8] that a certain statement that is related to Theorem 1.1 is equivalent to this conjecture. Within this context (and others), Theorem 1.1 has been studied, reproved, and generalized many times including results by Vershynin in [21], by Spielman and Srivastava in [20] (who showed that for $0 < \varepsilon < 1$ one can choose $c_0 = \varepsilon^2$ and $d_0 = (1 - \varepsilon)$), and by Naor and Youssef in [18]. Using some of the techniques developed in [20], Marcus, Spielman, and Srivastava eventually solved the Kadison-Singer conjecture in [16].

The preservation of lower ℓ_2 -norm bounds on subspaces is also related to a problem from infinite-dimensional Banach space theory, namely the factorization property of bounded linear operators with large diagonal. This problem has its origins in Andrew’s paper [3] and it has been further developed by Laustsen, Lechner, and Müller in [11], by Lechner in [12], by Lechner, Müller, Schlumprecht, and the third author in [13] and [14], and others. In the finite-dimensional Euclidean setting, the problem can be stated as follows: Given $n \in \mathbb{N}$ and $\theta > 0$, determine $C > 0$ and $m \in \mathbb{N}$ such that every norm-one $n \times n$ matrix $A = (a_{i,j})$ with $\min |a_{i,i}| \geq \theta$ is a C/θ -factor of the $m \times m$ identity matrix I_m . This means that there exist matrices L and R of appropriate dimension such that $\|L\|\|R\| \leq C/\theta$ and $LAR = I_m$. By (2) one may take $C = 2$ and $m \geq (3/4)n\theta^2$. The continuous version of this problem was investigated by Dai, Hore, Jiao, Lan, and the third author in [9] where non-optimal estimates were given.

We turn out attention towards formulating a version of Theorem 1.1 in the setting of a continuous matrix function $A : \mathbb{R} \rightarrow M_{n \times n}$. A point-wise application of Theorem 1.1 gives that there exists a choice of coordinate subspaces of constant dimension $U(t) \subset \mathbb{R}^n$, for all $t \in \mathbb{R}$ such that $v \in U(t)$, $\|A(t)v\| \geq c\|v\|$. However, the collection $\{U(t)\}_{t \in \mathbb{R}}$ is not a priori known to satisfy any useful properties. The property

of focus in this paper is continuity. We say that a collection $\{U(t)\}_{t \in \mathbb{R}}$ of subspaces varies continuously or is a continuous collection of subspaces if the matrix function $P: \mathbb{R} \rightarrow M_{n \times n}$, assigning to each $t \in \mathbb{R}$ the orthogonal projection $P(t)$ onto $U(t)$, is continuous (see Definition 2.3 and Theorem 2.5). We focus on generalizing Theorem 1.1 by formulating two problems.

PROBLEM 1. Given a continuous matrix function $A: \mathbb{R} \rightarrow M_{n \times n}(\mathbb{R})$ satisfying pointwise the same hypotheses as those in Theorem 1.1, is it possible to find subspaces $\{U(t)\}_{t \in \mathbb{R}}$ that vary continuously satisfying a similar lower ℓ_2 -norm bound? If so, are there obtainable bounds on $\dim(U(t))$?

Without involving the Bourgain-Tzafriri theorem we use elementary techniques from linear algebra and analysis to obtain the desired collection $\{U(t)\}_{t \in \mathbb{R}}$ with a lower bound of the dimension which is indeed optimal up the imposed restriction on the matrix function A and up to a universal constant. More precisely, in Section 3 we prove the following result, in which no structure for the subspaces $U(t)$ is guaranteed.

THEOREM 1.2. *Let $A: \mathbb{R} \rightarrow M_{n \times n}$ be a continuous matrix function, satisfying the property that for all $t \in \mathbb{R}$, $\|A(t)e_i\| = 1$ for every $1 \leq i \leq n$. Let $\Lambda = \sup_t \|A(t)\|$ and $\gamma \in (0, 1)$. Then, there exists a continuous family of m -dimensional subspaces $\{U(t)\}_{t \in \mathbb{R}}$ where $m \geq (1 - \gamma^2)n/(7\Lambda^2)$ such that for every $t \in \mathbb{R}$, and every $v \in U(t)$, $\|Av\| \geq \gamma\|v\|$.*

Note that the Hilbert-Schmidt norm of each $A(t)$ is \sqrt{n} and therefore $\Lambda \leq \sqrt{n}$.

If we wish to obtain a more rigid collection of subspaces $\{U(t)\}_{t \in \mathbb{R}}$, and thus a statement that is more similar to the Bourgain-Tzafriri theorem, then we must reformulate the problem. To that end we introduce the following notion, that is compatible to continuously varying subspaces.

DEFINITION 1.3. An m -dimensional subspace U of \mathbb{R}^n is called a quadratic convex combination of disjoint basis vectors if there exist disjoint subsets $\sigma_1 = \{i_1 < \dots < i_m\}$, $\sigma_2 = \{j_1 < \dots < j_m\}$ of $\{1, \dots, n\}$ and $\lambda \in [0, 1]$ such that U is spanned by the orthonormal sequence $u_k = \lambda^{1/2}e_{i_k} + (1 - \lambda)^{1/2}e_{j_k}$, $k = 1, \dots, m$. If we wish to be more specific, we will say that U is a λ -quadratic convex combination of U_{σ_1} and U_{σ_2} .

PROBLEM 2. Given a continuous matrix function $A: \mathbb{R} \rightarrow M_{n \times n}(\mathbb{R})$ satisfying pointwise the same hypotheses as those in Theorem 1.1, is it possible to find subspaces $\{U(t)\}_{t \in \mathbb{R}}$ that vary continuously, satisfy a similar lower ℓ_2 -norm bound, and such that for all $t \in \mathbb{R}$, $U(t)$ is a quadratic convex combinations of disjoint basis vectors? If so, are there obtainable bounds on $\dim(U(t))$?

Following the original proof of Bourgain and Tzafriri from [5], we deploy probabilistic and combinatorial techniques to give an answer to Problem 2. As a trade-off for the structural properties of the obtained continuous choice of subspaces, the dimensional estimate is weakened. The following, is the main result of our paper.

THEOREM 1.4. *There exists universal constants $c, d > 0$ such that for all continuous matrix functions $A : \mathbb{R} \rightarrow M_{n \times n}$ with the property that $\|A(t)e_i\| = 1$ for all $t \in \mathbb{R}$ and $1 \leq i \leq n$, there exists a continuous family of m -dimensional subspaces $\{U(t)\}_{t \in \mathbb{R}}$ of \mathbb{R}^n with $m \geq dn/\Lambda^4$ where $\Lambda = \sup_t \|A(t)\|$ such that $\|Av\| \geq c\|v\|$ for every $t \in \mathbb{R}$ and every $v \in U(t)$. Furthermore, each subspace $U(t)$ is a quadratic convex combination of disjoint basis vectors.*

The impact of this improved choice of subspaces on the dimensional estimate is on the exponent of Λ . In light of Theorem 1.2, it is important to justify why such a steep price in dimension must be paid in the second method, which is presented in Section 4. We begin with a consideration of the issues that arise with a pointwise application of the Bourgain-Tzafriri result. Were we to try to merely apply Theorem 1.1 pointwise given $A : \mathbb{R} \rightarrow M_{n \times n}$, we would be able to generate a suitable collection $\{\sigma_t\}_{t \in \mathbb{R}}$ of subsets of $\{1, \dots, n\}$. Of course, this is not enough: U_{σ_t} cannot be chosen uniformly with respect to t , so there is no reason to suppose that for two arbitrarily close points t_1, t_2 the corresponding $U_{\sigma_{t_1}}, U_{\sigma_{t_2}}$ would serendipitously satisfy the required continuous transition property. To see this, it is enough to notice that satisfactory sets σ_t at a point need not be unique.

Instead we find a suitable countable sequence of points $(t_i)_{i \in \mathbb{Z}}$, apply a static result à la Bourgain-Tzafriri at each such t_i to first find a subset σ_i of $\{1, \dots, n\}$ and then conclude by means of a “stitching” argument. By this, we mean that we continuously pass between U_{σ_i} and $U_{\sigma_{i+1}}$ on $[t_i, t_{i+1}]$ via quadratic convex combinations $U(t)$ of U_{σ_i} and $U_{\sigma_{i+1}}$ while simultaneously preserving the desired lower ℓ_2 -norm bound of $A(t)$ on $U(t)$. Importantly, it is necessary that on the whole interval $[t_i, t_{i+1}]$, $A(t)$ already satisfies this bound on both U_{σ_i} and $U_{\sigma_{i+1}}$. But this alone is not sufficient. Without further stipulations, $U(t)$ could fail to preserve the desired properties at some point (e.g., $U(t)$ and $\ker A(t)$ may momentarily intersect for some $t \in [t_i, t_{i+1}]$, which would immediately violate the lower ℓ_2 -norm bound). From this, it becomes clear that there must be some relationship between σ_i and σ_{i+1} that would preserve the minimal stretch property during the “stitching” procedure. It is sufficient to require that for all $t \in [t_i, t_{i+1}]$, $A(t)$ satisfies the desired bound on the whole subspace $U_{\sigma_i \cup \sigma_{i+1}}$.

To achieve this, we rely on an iterative application of a modified Bourgain-Tzafriri argument. The modification introduces a dependence of each subsequent column set on the one that precedes it; the iteration consists of passing to subsets of the generated column sets, and allows for the modification to be bilateral. Then, each column set depends on both the one that precedes it and on the one that will follow. This is where the loss of optimality occurs.

Because the iterative application of the modified Bourgain-Tzafriri result incurs the loss of optimal dimension, in the proof of Theorem 1.2, we base the argument on a different way of refining the column sets. We introduce a dependence that allows for a different “stitching” argument to be made which takes place at the level of the spaces $A(t)(U_{\sigma_i})$ and $A(t)(U_{\sigma_{i+1}})$. This is based on nice properties of orthogonal subspaces, to which one may conveniently pass with a judicious construction. As the spectral decomposition estimate is applied once and subsequent modifications come only at the cost of worse universal constants, this optimally solves Problem 1.

Closely related to Problem 1, is the following problem considered in [9].

PROBLEM 3. Given $n \in \mathbb{N}$ and $\theta > 0$, determine $C > 0$ and $m \in \mathbb{N}$ such that for every continuous matrix function $A = (a_{i,j}) : \mathbb{R} \rightarrow M_{n \times n}$ with $\|A(t)\| \leq 1$ and $\min_i |a_{i,i}(t)| \geq \theta$ for every $t \in \mathbb{R}$ there exists continuous matrix functions $L : \mathbb{R} \rightarrow M_{m \times n}$, $R : \mathbb{R} \rightarrow M_{n \times m}$ so that $L(t)A(t)R(t) = I_m$ and $\sup_t \|L(t)\| \|R(t)\| \leq C/\theta$.

In Theorem 3.10 we will show that a solution to Problem 1 also offers one for Problem 3.

The paper is organized as follows. Section 2 recalls the preliminaries of matrices and continuous matrix functions. Section 3 is devoted to the proof of Theorem 1.2 using elementary tools. The expert reader may wish to skip directly to Section 4, in which we prove Theorem 1.4 by adapting the original approach of Bourgain and Tzafriri from [5]. In Section 5 we briefly discuss possible future directions of this line of research.

2. Preliminaries

In this section we recall various norms, recall the concept of continuous matrix functions, and discuss continuously varying subspaces of \mathbb{R}^n . We denote the standard basis for \mathbb{R}^n by $\{e_1, \dots, e_n\}$ and assume the norm on \mathbb{R}^n to be the 2-norm: $\|x\| = (\sum_{i=1}^n x_i^2)^{1/2}$ for all $x \in \mathbb{R}^n$. We begin by defining useful quantities on $M_{m \times n}$, the space of all real valued $m \times n$ matrices.

DEFINITION 2.1. Let $A = (a_{i,j}) \in M_{m \times n}$.

(i) The operator norm of A is defined as

$$\|A\| = \sup \left\{ \|Ax\| : x \in \mathbb{R}^n, \|x\| = 1 \right\}.$$

If U is a subspace of \mathbb{R}^n , let $\|A|_U\| = \sup \{ \|Ax\| : x \in U, \|x\| = 1 \}$.

(ii) The minimal stretch of A is defined as

$$m_A = \inf \left\{ \|Ax\| : x \in \mathbb{R}^n, \|x\| = 1 \right\}.$$

If U is a subspace of \mathbb{R}^n , let $m_{A|_U} = \inf \{ \|Ax\| : x \in U, \|x\| = 1 \}$.

(iii) The Hilbert-Schmidt norm of A is defined as

$$\|A\|_{\text{HS}} = \left(\sum_{i=1}^n \|Ae_i\|^2 \right)^{1/2} = \left(\sum_{i=1}^m \sum_{j=1}^n a_{ij}^2 \right)^{1/2}.$$

We point out that these quantities can be equivalently defined as follows. Let $(\sigma_i)_{i=1}^{m \wedge n}$ denote sequence of the singular values of A . Then $\|A\| = \max \sigma_i$ whereas

$m_A = \min \sigma_i$ and $\|A\|_{\text{HS}} = (\text{tr}(A^T A))^{1/2} = (\sum_{i=1}^{m \wedge n} \sigma_i^2)^{1/2}$ (see, e.g., [15, Theorem 7.4.3]). These yield the following estimates.

$$\|A\| \leq \|A\|_{\text{HS}} \leq (\text{rank}(A))^{1/2} \|A\|.$$

For an $n \times n$ matrix the formula $\|A\|_{\text{HS}} = (\sum_{i=1}^n \sigma_i^2)^{1/2}$ is also used to deduce (2). Take a singular value decomposition $A = W \Sigma V^T$, where $\Sigma = \text{diag}(\sigma_1, \dots, \sigma_n)$ is the singular value matrix of A and $W, V = [v_1 \cdots v_n]$ are orthogonal matrices. Then, for any $\gamma \in (0, 1)$ the space $U = \langle \{v_i : \sigma_i \geq \gamma\} \rangle$ satisfies the property that for all $x \in U$, $\|Ax\| \geq \gamma \|x\|$. Set $E = \{v_i : \sigma_i \geq \gamma\}$ and observe

$$\|A\|_{\text{HS}}^2 = \sum_{i \in E} \sigma_i^2 + \sum_{i \notin E} \sigma_i^2 < \#E \|A\|^2 + \gamma^2 n.$$

Solving for $\#E$ yields (2).

DEFINITION 2.2. Let I be an interval of \mathbb{R} . A matrix function $A = (a_{i,j}) : I \rightarrow M_{m \times n}$ is said to be continuous if its entries $a_{i,j} : I \rightarrow M_{m \times n}$ are continuous.

It is well known that A is continuous if and only if it is continuous as a function from the metric space $(I, |\cdot|)$ to the normed linear space $(M_{m \times n}, \|\cdot\|)$ (see, e.g., [9, Lemma 3.1]).

DEFINITION 2.3. Let I be an interval of \mathbb{R} and let $\{U(t)\}_{t \in I}$ be a family of subspaces of \mathbb{R}^n .

- (i) The family $\{U(t)\}_{t \in I}$ is said to be a continuous choice of subspaces of \mathbb{R}^n if the function $P : I \rightarrow M_{n \times n}$ where $P(t)$ is the orthogonal projection onto $U(t)$ for all $t \in I$ is continuous.
- (ii) The family $\{U(t)\}_{t \in I}$ is said to admit a continuous choice of basis if, for some $m \in \mathbb{N}$, there exist continuous $\gamma_1, \dots, \gamma_m : I \rightarrow \mathbb{R}^n$ so that for each $t \in I$, $\gamma_1(t), \dots, \gamma_m(t)$ form a basis of $U(t)$.

LEMMA 2.4. If I is an interval of \mathbb{R} and a family $\{U(t)\}_{t \in I}$ of subspaces of \mathbb{R}^n admits a continuous choice of basis $\gamma_1, \dots, \gamma_m : I \rightarrow \mathbb{R}^n$ then it also admits a continuous choice of orthonormal basis $u_1, \dots, u_m : I \rightarrow \mathbb{R}^n$. Therefore $\{U(t)\}_{t \in I}$ is a continuous choice of subspaces of \mathbb{R}^n .

Proof. Define $u_1(t) = \|\gamma_1(t)\|^{-1} \gamma_1(t)$. By induction on $k = 2, \dots, m$ let $\tilde{u}_k(t) = \gamma_k(t) - \sum_{i=1}^{k-1} \langle \gamma_k(t), u_i(t) \rangle u_i(t)$ and $u_k(t) = \|\tilde{u}_k(t)\|^{-1} \tilde{u}_k(t)$. This yields a continuous choice of orthonormal basis $u_1, \dots, u_m : I \rightarrow \mathbb{R}^n$. Finally, note that for each $t \in I$, putting $W(t) = [u_1(t) \cdots u_m(t)]$ we have that $P(t) = W(t)W^T(t)$ the orthogonal projection onto $U(t)$ and $P : I \rightarrow M_{n \times n}$ is continuous. \square

Our next goal is to show that (i) and (ii) of Definition 2.3 are in fact equivalent. Note it is essential that the domain I is a subset of \mathbb{R} . Indeed if we replace I with the

unit circle S^1 then this would no longer be true since we are working with subspaces of \mathbb{R}^n . For example, for all $x = (\cos(\theta), \sin(\theta))$ in S^1 let

$$U(\cos(\theta), \sin(\theta)) = \langle (\cos(\theta/2), \sin(\theta/2)) \rangle \subset \mathbb{R}^2.$$

Then $\{U(x)\}_{x \in S^1}$ is a continuous choice of subspaces of \mathbb{R}^2 for which a continuous choice of basis is impossible.

THEOREM 2.5. *Let I be an interval of \mathbb{R} . A family of subspaces $\{U(t)\}_{t \in I}$ of \mathbb{R}^n is continuous if and only if it admits a continuous choice of orthonormal basis.*

The proof of the above requires some preparatory steps. For each of the following lemmata, let I be an interval of \mathbb{R} and $P : I \rightarrow M_{n \times n}$ be a continuous matrix function such that $P(t)$ is an orthogonal projection for all $t \in \mathbb{R}$.

LEMMA 2.6. *The rank of $P(t)$ is constant for all $t \in I$.*

Proof. Since the rank of an orthogonal projection is equal to the trace of the orthogonal projection, and since $\text{tr}(P(t)) : I \rightarrow \mathbb{R}$ is continuous when P is continuous, the result follows. \square

LEMMA 2.7. *For all $t_0 \in I$ there is an $\varepsilon > 0$ and continuous functions $\gamma_1, \dots, \gamma_k : (t_0 - \varepsilon, t_0 + \varepsilon) \cap I \rightarrow \mathbb{R}^n$ such that for all $t \in (t_0 - \varepsilon, t_0 + \varepsilon) \cap I$, $\mathfrak{Z}(P(t)) = \langle \gamma_1(t), \dots, \gamma_k(t) \rangle$.*

Proof. Let $t_0 \in I$. By assumption, $\text{rank}(P(t_0)) = k$, so there exist k linearly independent columns of $P(t_0)$, $\{Pe_{i_j}(t_0)\}_{j=1}^k$, which span $\mathfrak{Z}(P(t_0))$. That is,

$$\mathfrak{Z}(P(t_0)) = \langle Pe_{i_1}(t_0) \dots Pe_{i_k}(t_0) \rangle.$$

Since $\text{rank}(Pe_{i_1}(t_0) \dots Pe_{i_k}(t_0)) = k$, we can find k suitable rows of the matrix $[Pe_{i_1}(t) \dots Pe_{i_k}(t)]$ to obtain $B : \mathbb{R} \rightarrow M_{k \times k}$ such that $B(t_0)$ is invertible (by virtue of satisfying $\det B(t_0) \neq 0$). The continuity of the determinant function guarantees the existence of an $\varepsilon > 0$ such that for any $t \in (t_0 - \varepsilon, t_0 + \varepsilon) \cap I$, $\det B(t) \neq 0$. Thus, for any $t \in (t_0 - \varepsilon, t_0 + \varepsilon) \cap I$, $\text{rank}(Pe_{i_1}(t) \dots Pe_{i_k}(t)) = k$ and so

$$\mathfrak{Z}(P(t)) = \langle Pe_{i_1}(t) \dots Pe_{i_k}(t) \rangle \text{ as } \text{rank}(P(t)) = k$$

(where $B(t)$ is the minor of $P(t)$ defined using the same components as $B(t_0)$). \square

LEMMA 2.8. *Let $a < c < b < d$ and $P : (a, d) \rightarrow M_{n \times n}$ be continuous and let $k = \text{rank}(P(t))$ for all $t \in \mathbb{R}$. Assume that $\mathfrak{Z}(P(t))$ admits a continuous choice of basis $\gamma_1, \dots, \gamma_k : (a, b) \rightarrow \mathbb{R}^n$ on (a, b) and another continuous choice of basis on (c, d) . Then, $\gamma_1, \dots, \gamma_k$ may be extended to a continuous choice of basis on (a, d) .*

Proof. By Lemma 2.7, it is sufficient to show that we can “stitch together” continuous bases with intersecting domains. Let

$$\begin{aligned}\alpha_1, \dots, \alpha_k &: (a, b) \rightarrow \mathbb{R}^n \text{ s.t. } \forall t \in (a, b), \operatorname{Im}(P(t)) = \langle \alpha_1(t), \dots, \alpha_k(t) \rangle \\ \beta_1, \dots, \beta_k &: (c, d) \rightarrow \mathbb{R}^n \text{ s.t. } \forall t \in (c, d), \operatorname{Im}(P(t)) = \langle \beta_1(t), \dots, \beta_k(t) \rangle\end{aligned}$$

for some $a < c < b < d$. Let $t_0 \in (c, b)$. Then we can write the $\alpha_i(t_0)$ as linear combinations of $\beta_1(t_0), \dots, \beta_k(t_0)$ since $\langle \alpha_1(t_0), \dots, \alpha_k(t_0) \rangle = \langle \beta_1(t_0), \dots, \beta_k(t_0) \rangle$. In other words, for each $i \in \{1, \dots, k\}$, there are $(\lambda_{ij})_{j=1}^k$ s.t. $\alpha_i(t_0) = \sum_{j=1}^k \lambda_{ij} \beta_j(t_0)$.

Since the $(\alpha_i(t_0))_{i=1}^k$ and $(\beta_i(t_0))_{i=1}^k$ are bases of $\operatorname{Im}(P(t_0))$, the λ_{ij} uniquely determine an invertible matrix $F = (\lambda_{ij}) \in M_{k \times k}(\mathbb{R})$, for which

$$\begin{bmatrix} \alpha_1(t_0)^T \\ \vdots \\ \alpha_k(t_0)^T \end{bmatrix} = F \begin{bmatrix} \beta_1(t_0)^T \\ \vdots \\ \beta_k(t_0)^T \end{bmatrix}.$$

Define $(\gamma_1, \dots, \gamma_k) : (a, d) \rightarrow M_{n \times k}(\mathbb{R})$ by:

$$(\gamma_1(t), \dots, \gamma_k(t)) = \begin{cases} (\alpha_1(t), \dots, \alpha_k(t)) & t \leq t_0 \\ (\beta_1(t), \dots, \beta_k(t)) F^T & t > t_0. \end{cases}$$

Then for every $t \in (a, d)$, $\operatorname{Im}(P(t)) = \langle \gamma_1(t), \dots, \gamma_k(t) \rangle$ (since $\det F \neq 0$), and the γ_i are continuous. \square

Proof of Theorem 2.5. Let $P : I \rightarrow M_{n \times n}$ be continuous such that for all $t \in I$, $P(t)$ is an orthogonal projection. We want to construct $u_i : I \rightarrow \mathbb{R}^n$ continuous such that $u_1(t), \dots, u_k(t)$ form a basis of $\operatorname{Im}(P(t))$ for every $t \in I$. For simplicity, let us assume that $I = \mathbb{R}$ as the other cases are similar.

We work first with $[-1, 1]$. As it is compact, by Lemma 2.7, $[-1, 1]$ admits a finite cover by open intervals (a_j, b_j) so that there exist continuous $\gamma_{ji} : (a_j, b_j) \rightarrow \mathbb{R}^n$ such that for $t \in (a_j, b_j)$, $(\gamma_{j1}(t), \dots, \gamma_{jk}(t))$ is a basis of $\operatorname{Im}(P(t))$.

Without loss of generality, we assume $a_{j+1} < b_j$ to obtain a cover by “interlocking” intervals (with non-empty sequential intersections). A finite, step-by-step application of Lemma 2.8 allows for the construction of continuous $\gamma_i : [-1, 1] \rightarrow \mathbb{R}^n$ such that $(\gamma_1(t), \dots, \gamma_k(t))$ is a basis of $\operatorname{Im}(P(t))$ for each $t \in [-1, 1]$.

Suppose $\gamma_i(t)$ is defined for all $t \in [-m, m]$. We will extend γ_i to $[-(m+1), m+1]$. We similarly use Lemma 2.7 and the compactness of $[-(m+1), m+1]$ to obtain a suitable finite open cover, whereupon we apply Lemma 2.8 as above to define $\gamma_i(t)$ for $t \in [-(m+1), m+1]$ without modifying γ_i on $[-m, m]$. Notice that γ_i is well-defined as it is independent of the choice of m (subsequent extensions of γ_i do not alter the behaviour of the function on a domain on which it was previously defined). \square

3. Method I: Optimal Dimensional Bound

In this section we prove Theorem 1.2 and then show how it can be used to provide a solution to Problem 3.

3.1. The proof of Theorem 1.2

We start by proving some statements that will allow us to find appropriate subspaces of \mathbb{R}^n and then stitch them together. Although this proof combines well known concepts, we have chosen to present some of their proofs for the purpose of making the paper as self contained as possible.

PROPOSITION 3.1. *If X and Y are linear subspaces of \mathbb{R}^n and $\dim X = \dim Y = m$, then there exist linear subspaces $\tilde{X} \subset X, \tilde{Y} \subset Y$ such that $\dim \tilde{X}, \dim \tilde{Y} \geq \lfloor \frac{m}{2} \rfloor$ and $\tilde{X} \perp \tilde{Y}$.*

We will prove Proposition 3.1 by using the notion of principal angles of subspaces (see, e.g., [1]), as it will also be used later.

LEMMA 3.2. *Given two subspaces X and Y of \mathbb{R}^n of dimension m , there exist orthonormal bases $\{x_1, x_2, \dots, x_m\}$ and $\{y_1, y_2, \dots, y_m\}$ of X and Y respectively, such that $\langle x_i, y_j \rangle = 0$ for any $i \neq j$.*

Proof. Define $P : \mathbb{R}^n \rightarrow Y$ to be the projection map onto Y . Let $\tilde{P} = P|_X$, and so, $\tilde{P} : X \rightarrow Y$. Define $A = \tilde{P}^* \tilde{P} : X \rightarrow X$. Then A is self-adjoint, so the Spectral Theorem implies there is an orthonormal basis $\{x_1, \dots, x_m\}$ of X such that each x_i is an eigenvector of A . Now, let $\tilde{y}_i = \tilde{P}x_i$ and $\sigma = \{1 \leq i \leq m : \tilde{y}_i \neq 0\}$. For $i \in \sigma$ let $y_i = \|\tilde{y}_i\|^{-1} \tilde{y}_i$. Note for $i, j \in \sigma$ with $i \neq j$ that

$$\langle \tilde{y}_i, \tilde{y}_j \rangle = \langle \tilde{P}x_i, \tilde{P}x_j \rangle = \langle x_i, \tilde{P}^* \tilde{P}x_j \rangle = \langle x_i, Ax_j \rangle = \lambda_j \langle x_i, x_j \rangle = 0$$

so the sequence $(y_i)_{i \in \sigma}$ is orthonormal. Extend $(y_i)_{i \in \sigma}$ to an orthonormal basis $(y_i)_{i=1}^m$ of Y . It now remains to check that $\langle x_i, y_j \rangle = 0$ for $i \neq j$. If $\tilde{P}x_i = 0$, then x_i is orthogonal to every vector in Y . Otherwise, notice that $u_i \in \mathbb{R}^n = Y \oplus Y^\perp$, and so we can express it as $x_i = \tilde{y}_i + y'_i$ where $y'_i = x_i - \tilde{y}_i \in Y^\perp$. Then, it follows:

$$\langle x_i, y_j \rangle = \langle \tilde{y}_i + y'_i, y_j \rangle = \langle \tilde{y}_i, y_j \rangle + \langle y'_i, y_j \rangle = 0. \quad \square$$

We can now proceed with the proof of Proposition 3.1

Proof. Lemma 3.2 yields orthonormal bases $\{x_1, \dots, x_m\}$ and $\{y_1, \dots, y_m\}$ of X and Y respectively, such that $\langle x_i, y_j \rangle = 0$ when $i \neq j$. When m is odd, define $\tilde{X} = \langle \{x_1, \dots, x_{\lfloor \frac{m}{2} \rfloor} \} \rangle$ and $\tilde{Y} = \langle \{y_{\lceil \frac{m}{2} \rceil}, \dots, y_{m-1}, y_m\} \rangle$. Then $\langle x_i, y_j \rangle = 0$ for all $1 \leq i \leq \lfloor \frac{m}{2} \rfloor, \lceil \frac{m}{2} \rceil \leq j \leq m$.

If m is even, define $\tilde{X} = \langle \{x_1, \dots, x_{\frac{m}{2}} \} \rangle$ and $\tilde{Y} = \langle \{y_{\frac{m}{2}+1}, \dots, y_{m-1}, y_m\} \rangle$. Then $\langle x_i, y_j \rangle = 0$ for all $1 \leq i \leq \frac{m}{2}, \frac{m}{2} + 1 \leq j \leq m$. Thus, $\tilde{X} \perp \tilde{Y}$. \square

By Proposition 3.1, given any two subspaces, we can always find orthogonal subspaces at the cost of reducing the dimension by half. The following result again appeals to Lemma 3.2 in order to show that given two subspaces of a fixed dimension, it is possible to traverse from one subspace to another through a continuously varying choice of subspaces of the same dimension. Note that this is also a standard argument used, e.g., to show the path-connectedness of the Grassmannian.

PROPOSITION 3.3. *Let $X, Y \subset \mathbb{R}^n$, with $\dim(X) = \dim(Y) = m$. Then, for any $a < b \in \mathbb{R}$, there exists a continuous choice of subspaces $\{U(t)\}_{t \in [a,b]}$ of \mathbb{R}^n , such that $X = U(a)$ and $Y = U(b)$, and $U(t)$ lies in $X + Y$ for all $t \in [a, b]$.*

Proof. Using Lemma 3.2, we can find an orthonormal bases $\{x_1, \dots, x_m\}$ and $\{y_1, \dots, y_m\}$ of X and Y respectively such that $\langle x_i, y_j \rangle = 0$ whenever $i \neq j$. Now define for $1 \leq i \leq m$,

$$u_i(t) = \begin{cases} \left(1 - \frac{t-a}{b-a}\right)x_i + \left(\frac{t-a}{b-a}\right)y_i & \text{if } x_i, y_i \text{ are linearly independent} \\ y_i & \text{if } x_i, y_i \text{ are linearly dependent.} \end{cases}$$

Let $U(t) = \langle u_1(t), u_2(t), \dots, u_m(t) \rangle$. Observe that $U(a) = X$ and $U(b) = Y$. Notice that for any $t \in [a, b]$, at no point do we have linear dependence between the u_i 's as, in fact, they are always orthogonal. Thus, $\{U(t)\}_{t \in [a,b]}$ as defined above is our required family of subspaces. \square

The following lemma guarantees the existence of a discrete collection of points in \mathbb{R}^n such that the application of Theorem 1.1 on all points of this collection preserves the desired minimal stretch property on overlapping intervals covering \mathbb{R} .

LEMMA 3.4. *Let $\varepsilon > 0$ and $A : \mathbb{R} \rightarrow M_{n \times n}(\mathbb{R})$ be a continuous matrix function. Then there exists a sequence $(t_i)_{i \in \mathbb{Z}}$ in \mathbb{R} such that the following are satisfied.*

(i) *For all $i \in \mathbb{Z}$, $t_i < t_{i+1}$, $\sup_i t_i = \infty$, and $\inf_i t_i = -\infty$.*

(ii) *For all $i \in \mathbb{Z}$ and t in $[t_{i-1}, t_{i+1}]$, $\|A(t_i) - A(t)\| \leq \varepsilon$.*

In particular, if $i \in \mathbb{Z}$, $c > 0$, and U is a subspace of \mathbb{R}^n such that $m_{A(t_i)|U} \geq c$ then for all $t \in [t_{i-1}, t_{i+1}]$, $m_{A(t)|U} \geq c - \varepsilon$.

Proof. By the uniform continuity of A on each compact interval $[i, i+1]$, there exists $k_i \in \mathbb{N}$ such that for all s, t in $[i, i+1]$ with $|s-t| \leq 1/k_i$, $\|A(s) - A(t)\| \leq \varepsilon/2$. Denote $F_i = \{i + j/k_i : j = 0, 1, \dots, k_i - 1\}$ and define $F = \cup_{i \in \mathbb{Z}} F_i$. Clearly, this is a discrete subset of \mathbb{R} that is unbounded above and unbounded below. Enumerating it in increasing order yields the desired sequence $(t_i)_{i \in \mathbb{Z}}$ that satisfies (i) and (ii).

For the final part let $i \in \mathbb{Z}$, $c > 0$, and U be a subspace of \mathbb{R}^n such that $m_{A(t_i)|U} \geq c$. Then, for all $x \in U$ with $\|x\| = 1$ and $t \in [t_{i-1}, t_{i+1}]$,

$$\|A(t)x\| \geq \|A(t_i)x\| - \|A(t_i) - A(t)\|\|x\| \geq c - \varepsilon.$$

In other words, $m_{A(t)|U} \geq c - \varepsilon$. \square

We can now prove Theorem 1.2, which we restate for convenience.

THEOREM 3.5. *Let $A : \mathbb{R} \rightarrow M_{n \times n}$ be a continuous matrix function, satisfying the property that for all $t \in \mathbb{R}$, $\|A(t)e_i\| = 1$ for every $1 \leq i \leq n$. Let $\Lambda = \sup_t \|A(t)\|$ and $\gamma \in (0, 1)$. Then, there exists a continuous family of m -dimensional subspaces $\{U(t)\}_{t \in \mathbb{R}}$ where $m \geq (1 - \gamma^2)n/(7\Lambda^2)$ such that for every $t \in \mathbb{R}$, and every $v \in U(t)$, $\|Av\| \geq \gamma\|v\|$.*


Proof. To begin, note it is always possible to find a continuous choice of one-dimensional subspaces by taking $U(t) = \langle e_1 \rangle$, for all $t \in \mathbb{R}$. Let $m_0 = \lceil (1 - \gamma^2)n/\Lambda^2 \rceil$. We will show that there is continuous choice of $\lfloor m_0/4 \rfloor$ -dimensional subspaces that satisfies the conclusion. Therefore, we can achieve $1 \vee \lfloor m_0/4 \rfloor$ which dominates $m_0/7$ for all possible values of m_0 .

Apply Lemma 3.4 to A for $0 < \varepsilon < (1 - \gamma)/2$ to find an increasing sequence of points $(t_i)_{i \in \mathbb{Z}}$ satisfying the conclusion of that lemma. As we will see later, ε may need to be smaller. We now argue that we can find subspaces U_i , $i \in \mathbb{Z}$, of common dimension m_0 such that for all $i \in \mathbb{Z}$ and $t \in [t_{i-1}, t_{i+1}]$ we have $m_{A(t)|U_i} \geq \gamma + \varepsilon$.

Let $\tilde{\gamma} = \gamma + 2\varepsilon < 1$ and for each $i \in \mathbb{Z}$ apply (2) to $A(t_i)$ to obtain a subspace U_i of common dimension $\tilde{m}_0 > (1 - \tilde{\gamma}^2)n/\Lambda^2$ such that for all $t \in [t_{i-1}, t_{i+1}]$, $m_{A(t)|U_i} \geq \tilde{\gamma} - \varepsilon = \gamma + \varepsilon$. As $\tilde{m}_0 \geq \lfloor (1 - \tilde{\gamma}^2)n/\Lambda^2 \rfloor + 1 = \lfloor (1 - (\gamma + 2\varepsilon)^2)n/\Lambda^2 \rfloor + 1$, for ε sufficiently small, we have $\tilde{m}_0 \geq \lceil (1 - \gamma^2)n/\Lambda^2 \rceil = m_0$. Later, ε may need to be made even smaller.

Let $i \in \mathbb{Z}$ be given. We will outline the mechanism by which one produces the required subspaces on $[t_i, t_{i+1}]$, and then conclude by extending the construction to all of \mathbb{R} by working on each interval and “stitching” together at the boundaries. Given i and $t \in [t_{i-1}, t_{i+1}]$ let $V_{i,t} := \langle A(t)e_j : j \in \sigma_i \rangle = A(t)(U_i)$.

A diagram illustrating the choice of subspaces on the interval $[t_i, t_{i+2}]$ is included below for the reader’s convenience.



Fix an $s_i \in (t_i, t_{i+1})$. At s_i , we apply Proposition 3.1 to the subspaces V_{i,s_i} and V_{i+1,s_i} to obtain subspaces $\tilde{V}_{i,s_i}^L \subset V_{i,s_i}$ and $\tilde{V}_{i+1,s_i}^R \subset V_{i+1,s_i}$ such that $\tilde{V}_{i,s_i}^L \perp \tilde{V}_{i+1,s_i}^R$ and $\dim \tilde{V}_{i,s_i}^L = \dim \tilde{V}_{i+1,s_i}^R = \lfloor \frac{m_0}{2} \rfloor$.

We once more invoke Proposition 3.1 at the pre-images of \tilde{V}_{i+1,s_i}^R and \tilde{V}_{i,s_i}^L under A to obtain subspaces $\tilde{U}_i^L \subset (A(s_i)|_{U_i})^{-1} \left(\tilde{V}_{i+1,s_i}^R \right) \subset U_i$ and $\tilde{U}_{i+1}^R \subset (A(s_i)|_{U_{i+1}})^{-1} \left(\tilde{V}_{i,s_i}^L \right) \subset U_{i+1}$ such that $\tilde{U}_i^L \perp \tilde{U}_{i+1}^R$. The iterative application of Proposition 3.1 incurs a further loss of dimension: for all $t \in [t_i, t_{i+1}]$,

$$\dim \tilde{U}_{i,t}^L = \dim \tilde{U}_{i+1,t}^R = \left\lfloor \frac{1}{2} \left\lfloor \frac{m_0}{2} \right\rfloor \right\rfloor = \left\lfloor \frac{m_0}{4} \right\rfloor.$$

Now, we must find a way to continuously “stitch” \tilde{U}_i^L and \tilde{U}_{i+1}^R around s_i , and similarly between \tilde{U}_{i+1}^R and \tilde{U}_{i+1}^L to pass to the next interval. This second “stitch” will be necessary as there is no reason to suppose that $\tilde{U}_{i+1}^R = \tilde{U}_{i+1}^L$.

To this end, we pick some $\eta_i \in (0, \min\{\frac{s_i - t_i}{2}, \frac{t_{i+1} - s_i}{2}\})$. This will be the margin through at which the “stitching” occurs (that is, there will be two instances of “stitching together”: on $(s_i - \eta_i, s_i + \eta_i)$ and on $(t_{i+1} - \eta_i, t_{i+1}]$).

Observe that an application of Proposition 3.3 allows us to switch from \tilde{U}_i^L at $s_i - \eta_i$ to \tilde{U}_{i+1}^R at $s_i + \eta_i$, through subspaces of dimension m_0 without violating the minimal stretch property as the collection of subspaces are contained in $\tilde{U}_i^L \oplus \tilde{U}_{i+1}^R$. This will be verified in Lemma 3.6.

It therefore remains only to “stitch” together on $(t_{i+1} - \eta_i, t_{i+1}] \subsetneq (s_i, t_{i+1}]$. Notice that $\tilde{U}_{i+1}^R \subset U_{i+1}$, and $\tilde{U}_{i+1}^L \subset U_{i+1}$. As the minimal stretch property holds for all vectors in U_{i+1} , the minimal stretch property also holds in $\tilde{U}_{i+1}^R + \tilde{U}_{i+1}^L$. Thus, we may appeal to Proposition 3.3 once again, and find a collection of subspaces of dimension $\lfloor \frac{m_0}{4} \rfloor$ to traverse from \tilde{U}_{i+1}^R at $t_{i+1} - \eta_i$, to \tilde{U}_{i+1}^L at t_{i+1} .

To recapitulate, on each $[t_i, t_{i+1}]$, we take:

$$\begin{aligned} & \tilde{U}_i^L \text{ for } t \in [t_i, s_i - \eta_i], \\ & \text{“stitch” } \tilde{U}_i^L \text{ and } \tilde{U}_{i+1}^R \text{ for } t \in (s_i - \eta_i, s_i + \eta_i), \\ & \tilde{U}_{i+1}^R \text{ for } t \in [s_i + \eta_i, t_{i+1} - \eta_i], \\ & \text{“stitch” } \tilde{U}_{i+1}^R \text{ and } \tilde{U}_{i+1}^L \text{ for } t \in (t_{i+1} - \eta_i, t_{i+1}), \\ & \tilde{U}_{i+1,t}^L \text{ for } t = t_{i+1}. \quad \square \end{aligned}$$

LEMMA 3.6. *For any vector x lying in $\tilde{U}_i^L \oplus \tilde{U}_{i+1}^R$, $i \in \mathbb{Z}$, the vector x satisfies $\|A(t)x\| \geq \gamma\|x\|$, when $t \in [t_i, t_{i+1}]$.*

Proof. If $x \in \tilde{U}_i^L \oplus \tilde{U}_{i+1}^R$, then $x = f + g$, where $f \in \tilde{U}_i^L$ and $g \in \tilde{U}_{i+1}^R$. As \tilde{U}_i^L and \tilde{U}_{i+1}^R satisfy the minimal stretch property, we know that $\|A(t)f\| \geq (\gamma + \varepsilon)\|f\|$ for $t \in [t_{i-1}, t_{i+1}]$, and $\|A(t)g\| \geq (\gamma + \varepsilon)\|g\|$ for $t \in [t_i, t_{i+2}]$. Then for $t \in [t_{i-1}, t_{i+1}]$, we have

$$\begin{aligned} \|A(t)x\|^2 &= \|A(t)f\|^2 + \|A(t)g\|^2 + 2\langle A(t)f, A(t)g \rangle \\ &\geq (\gamma + \varepsilon)^2\|f\|^2 + (\gamma + \varepsilon)^2\|g\|^2 + 2\langle A(t)f, A(t)g \rangle \\ &= (\gamma + \varepsilon)^2\|x\|^2 + 2\langle A(t)f, A(t)g \rangle. \end{aligned} \tag{3}$$

Now recall that $A(s_i)f \in \tilde{V}_{i+1,s_i}^R$ and $A(s_i)g \in \tilde{V}_{i,s_i}^L$ so

$$\begin{aligned} |\langle A(t)f, A(t)g \rangle| &= |\langle A(t)f - A(s_i)f, A(t)g \rangle + \langle A(s_i)f, A(t)g - A(s_i)g \rangle + \langle A(s_i)f, A(s_i)g \rangle| \\ &\leq (\|A(t) - A(s_i)\| \|f\|) (\|A(t)\| \|g\|) \\ &\quad + (\|A(s_i)\| \|f\|) (\|A(t) - A(s_i)\| \|g\|) + 0 \\ &\leq \Lambda \|A(t) - A(s_i)\| \|x\|^2. \end{aligned}$$

From Lemma 3.4 (ii), we know that $\|A(t) - A(s_i)\| \leq \varepsilon$. Substituting this in (3), we obtain

$$\|A(t)x\|^2 \geq (\gamma + \varepsilon)^2 \left(1 - 2\Lambda\varepsilon(\gamma + \varepsilon)^{-2}\right) \|x\|^2.$$

By taking ε sufficiently small we obtain $\|A(t)x\| \geq \gamma \|x\|$. \square

REMARK 3.7. We sketch an argument that demonstrates that the dimensional estimate $n/\|A\|^2$ in Theorem 1.1 (and hence also in Theorem 1.2) is optimal, up to a constant. Assume that for $n \in \mathbb{N}$ and $1 \leq \lambda \leq \sqrt{n}$, there is a $m(n, \lambda) \in \mathbb{N}$ such that for any $n \times n$ matrix A with unit-length columns and $\|A\| \leq \lambda$, there exists $\sigma \subset \{1, \dots, n\}$ with $|\sigma| \geq m(n, \lambda)$ and $m_{A|_{U_\sigma}} > 0$. We will show that necessarily $m(n, \lambda) < 4n/\lambda^2$. To find an A that confirms this, take $m = \lceil n/\lambda^2 \rceil$ and using Euclidean division write $n = dm + r$. Find an orthonormal sequence u_1, \dots, u_m in \mathbb{R}^n such that $\langle u_1, \dots, u_m \rangle \perp \langle e_1, \dots, e_r \rangle$. Define $U = [u_1 \cdots u_m] \in M_{n \times m}$, $B = [U | \cdots | U] \in M_{n \times dm}$ (take d copies of U), $D = [e_1 \cdots e_r] \in M_{n \times r}$ and $A = [D | B] \in M_{n \times n}$. Then, A has unit-length columns, $\|A\| = \sqrt{d} \leq \lambda$, and $\text{rank}(A) = m + r < 2m \leq 4n/\lambda^2$. Therefore, for any $\sigma \subset \{1, \dots, n\}$ such that $m_{A|_{U_\sigma}} > 0$ must satisfy $m(n, \lambda) \leq |\sigma| \leq \text{rank}(A) < 4n/\lambda^2$.

REMARK 3.8. Although the dimensional estimate $n/\|A\|^2$ is optimal relative to the quantity $\|A\|$ and presupposing that A has unit-length columns. This does not preclude the existence of better dimensional estimates relative to other quantities related to a matrix A such as the stable rank, the p -stable rank, and the entropic stable rank of A . These quantities are based in the singular values of A and can be more useful in some settings (e.g., when studying matrices that don't necessarily have unit length columns). This topic has been studied, e.g., in [21], [20], [18], and [17].

3.2. Continuous factorization of the identity

In this subsection we explain how Theorem 1.2 yields a solution to Problem 3. We begin with a lemma that will help us find the left matrix in the factorization of the identity.

LEMMA 3.9. *Let I be an interval of \mathbb{R} and $A : I \rightarrow M_{n \times m}$ be a continuous matrix function such that there exists $c > 0$ with $m_{A(t)} \geq c$ for all t . Then there exists a continuous matrix function $L : \mathbb{R} \rightarrow M_{m \times n}$ such that $L(t)A(t) = I_m$ and $\|L(t)\| \leq 1/c$ for all $t \in I$.*

Proof. For each fixed $t \in I$ we have that $m_{A(t)} > 0$ and thus $A(t)$ has trivial kernel. This implies that the $m \times m$ matrix $A^T(t)A(t)$ has trivial kernel and is thus invertible. Since the matrix function $A^T A : I \rightarrow M_{m \times m}$ is continuous and pointwise invertible, $(A^T A)^{-1} : I \rightarrow M_{m \times m}$ is continuous as well (see, e.g., [9, Lemma 3.2]). Then, $L = (A^T A)^{-1} A^T : I \rightarrow M_{m \times n}$ is continuous and for each $t \in I$, $L(t)A(t) = I_m$. To find $\|L(t)\|$, for fixed $t \in I$, write the singular value decomposition $A(t) = U \Sigma V^T$ where U, V are unitary and Σ is rectangular diagonal. A direct computation yields $L(t) = V \tilde{\Sigma} U^T$, where $\tilde{\Sigma} = (\Sigma^T \Sigma)^{-1} \Sigma^T$, which is the matrix formed by taking reciprocals of all non-zero diagonal elements of Σ and then taking the transpose. Since V, U are unitary and thus preserve the matrix norm under multiplication, $\|L(t)\| = \|\tilde{\Sigma}\| \leq 1/c$. \square

To find the right matrix, we will use Theorem 1.2 and Theorem 2.5.

THEOREM 3.10. *Let $A : \mathbb{R} \rightarrow M_{n \times n}$ be a continuous matrix function and $\theta > 0$ with $\|A(t)e_i\| \geq \theta$ for $1 \leq i \leq n$ and $\|A(t)\| \leq 1$ for $t \in \mathbb{R}$. Then for any $\gamma \in (0, 1)$ and $m \leq \lceil (1 - \gamma^2)n\theta^2/7 \rceil$ there exist continuous matrix functions L, R of appropriate dimensions such that $L(t)A(t)R(t) = I_m$ with $\|L(t)\|\|R(t)\| \leq (\gamma\theta)^{-1}$ for all $t \in \mathbb{R}$.*

Proof. Let $a_k(t)$ denote the k th column of $A(t)$,

$$D(t) = \left[\frac{e_1}{\|a_1(t)\|} \cdots \frac{e_n}{\|a_n(t)\|} \right] \quad \text{and} \quad \tilde{A}(t) = A(t)D(t).$$

Note that

$$\|\tilde{A}(t)\| \leq \|A(t)\| \max_{1 \leq i \leq n} \|a_i(t)\|^{-1} \leq \frac{1}{\theta}.$$

Also, A satisfies the hypothesis of Theorem 3.5, so that we have the existence of a collection of m -dimensional subspaces $\{U(t)\}_{t \in \mathbb{R}}$ that vary continuously such that $m_{\tilde{A}(t)|_{U(t)}} \geq \gamma$ for all $t \in \mathbb{R}$, where

$$m \geq \frac{1 - \gamma^2}{7} \frac{n}{\sup_t \|\tilde{A}(t)\|^2} \geq \frac{1 - \gamma^2}{7} n \theta^2.$$

By Theorem 2.5 there exists a continuous choice of orthonormal basis $u_1, \dots, u_m : \mathbb{R} \rightarrow \mathbb{R}^n$. Define $W(t) = [u_1(t) \cdots u_m(t)]$, which is a continuous matrix function with the property $\text{Im}(W(t)) = U(t)$ and for all $x \in \mathbb{R}^m$, $\|W(t)x\| = \|x\|$. Let $R(t) = D(t)W(t)$, which is continuous and satisfies $\|R(t)\| = \|D(t)\| \leq 1/\theta$.

Now since

$$\begin{aligned} m_{A(t)R(t)} &= \inf \left\{ \|\tilde{A}(t)W(t)x\| : x \in \mathbb{R}^m, \|x\| = 1 \right\} \\ &= \inf \left\{ \|\tilde{A}(t)x\| : x \in U(t), \|x\| = 1 \right\} \geq \gamma \end{aligned}$$

we can apply Lemma 3.9 to $A(t)R(t)$ to show the existence of a continuous left inverse $L(t)$ of $A(t)R(t)$ that satisfies $\|L(t)\| \leq \gamma^{-1}$ for all $t \in \mathbb{R}$. \square

4. Method II: Column Space Approach

In this section, we prove Theorem 1.4. The goal is to attain a continuous choice of subspaces on which a given matrix function satisfies the desired minimal stretch property that more closely resembles spaces spanned by a subset of the unit vector basis. These will be quadratic convex combinations of disjoint basis vectors (see Definition 1.3). Therefore, we require a statement that guarantees the existence of suitable pairs of subspaces spanned by disjoint basis vectors that behave sufficiently well with one another so that they can be “stitched” together.

4.1. A Bourgain-Tzafriri Theorem for disjoint subsets of the basis

We first prove the static result that is necessary in the proof of Theorem 1.4.

THEOREM 4.1. *There exist constants $1 > d_1 > d_2 > d_3 > 0$ such that the following holds. For every $n \times n$ matrix A with $\|Ae_i\| = 1$ for $i = 1, \dots, n$, and for every $\sigma_1 \subset \{1, \dots, n\}$ with $|\sigma_1| \leq d_1 n \|A\|^{-2}$, there exists*

$$\sigma_2 \subset \{1, \dots, n\} \setminus \sigma_1 \text{ with } |\sigma_2| \geq d_2 n \|A\|^{-2}$$

such that for any choice of scalars $\{a_j\}_{j \in \sigma_1 \cup \sigma_2}$, we have

$$\left\| \sum_{j \in \sigma_1 \cup \sigma_2} a_j A e_j \right\| \geq \frac{1}{16\sqrt{2}} \left(\sum_{j \in \sigma_2} |a_j|^2 \right)^{1/2}.$$

If we additionally assume that $|\sigma_1| \geq d_2 n \|A\|^{-2}$ and $d_1 d_2 n \|A\|^{-4} \geq 1$ then there also exist

$$\tau_1 \subset \sigma_1 \text{ and } \tilde{\sigma}_2 \subset \sigma_2 \text{ with } |\tau_1| \geq d_3 n \|A\|^{-4} \text{ and } |\tilde{\sigma}_2| \geq d_3 n \|A\|^{-4}$$

such that for any choice of scalars $\{a_j\}_{j \in \tau_1 \cup \tilde{\sigma}_2}$, we have

$$\left\| \sum_{j \in \tau_1 \cup \tilde{\sigma}_2} a_j A e_j \right\| \geq \frac{1}{32} \left(\sum_{j \in \tau_1 \cup \tilde{\sigma}_2} |a_j|^2 \right)^{1/2}.$$

In order to make the theorem more tractable, the proof is broken down into lemmata that follow the general shape of the argument first made by Bourgain and Tzafriri in [5]. The main difference is that the selection is performed in the complement of a fixed subset of the index set. Lemma 4.3 follows the outline of [5, Lemma 1.4], Lemma 4.4 follows the outline of [5, Theorem 1.5] and Lemma 4.4 follows the outline of the final step of [5, proof of Theorem 1.2, page 145]. It is worth pointing out that Bourgain and Tzafriri in [5] offered two arguments for the final step of the proof of their main theorem. The first one is based on an exhaustion argument and Khintchine’s inequality. The second one uses a Maurey-Nikishin factorization argument that involves the little Grothendieck Theorem. We have opted to follow the more elementary first approach (see, e.g., also [4, Lemma B]).

In Lemmata 4.3, 4.4, and 4.5 below it is assumed that we are given an $n \times n$ matrix A that satisfies the assumptions of Theorem 4.1 (i.e., $\|Ae_i\| = 1$ for $i = 1, \dots, n$).

REMARK 4.2. The constants in the Theorem 4.1 are $d_1 = 1/256$ (proof of Lemma 4.3), $d_2 = d_1/4$ (Lemma 4.4), and $d_3 = d_2^2/2$ (proof of Lemma 4.5).

LEMMA 4.3. *There exists a constant $d_1 > 0$ such that the following holds: For every $\sigma_1 \subset \{1, \dots, n\}$ with $|\sigma_1| \leq d_1 n \|A\|^{-2}$ there exists $\sigma_2 \subset \{1, \dots, n\} \setminus \sigma_1$ with $|\sigma_2| \geq d_1 n \|A\|^{-2}$ such that for every $i \in \sigma_2$,*

$$\|P_{\langle Ae_j : j \in (\sigma_1 \cup \sigma_2) \setminus \{i\} \rangle} Ae_i\| < \frac{1}{\sqrt{2}}.$$

Proof. Take $\delta = 1/(8\|A\|^2)$ and $d_1 = 1/(8 \cdot 32)$. Let $\{\xi_i\}_{i \in \sigma_1^c}$ be a sequence of independent random variables of mean δ over a probability space (Ω, Σ, μ) taking only the values 0 and 1. For each $i \in \sigma_1^c$, $\text{Var}(\xi_i) = \delta(1 - \delta)$ and by independence we obtain $\text{Var}(\sum_{i \in \sigma_1^c} \xi_i) = |\sigma_1^c| \delta(1 - \delta)$. Define

$$D = \left\{ \omega \in \Omega : \left| \sum_{i \in \sigma_1^c} \xi_i(\omega) - \delta |\sigma_1^c| \right| \geq \delta |\sigma_1^c| / 2 \right\}.$$

By Chebyshev's inequality,

$$\mu(D) \leq 4 \frac{\text{Var}(\sum_{i \in \sigma_1^c} \xi_i)}{\delta^2 |\sigma_1^c|^2} \leq 4(\delta |\sigma_1^c|)^{-1}. \quad (4)$$

For each $\omega \in \Omega$, let

$$\sigma(\omega) = \{j \in \sigma_1^c : \xi_j(\omega) = 1\}.$$

We will show that there must be at least one $\sigma(\omega)$ which can be modified slightly to have the desired property. Set $V = \langle Ae_i : i \in \sigma_1 \rangle$. Then

$$\begin{aligned} & \int_{\Omega} \sum_{i \in \sigma_1^c} \xi_i(\omega) \left\| P_{\langle \xi_j(\omega) Ae_j \cup V \rangle_{j \neq i, j \in \sigma_1^c}} (Ae_i) \right\|^2 d\mu \\ &= \sum_{i \in \sigma_1^c} \left(\int_{\Omega} \xi_i(\omega) d\mu \right) \left(\int_{\Omega} \left\| P_{\langle \xi_j(\omega) Ae_j \cup V \rangle_{j \neq i, j \in \sigma_1^c}} (Ae_i) \right\|^2 d\mu \right) \\ &= \delta \int_{\Omega} \sum_{i \in \sigma_1^c} \left\| P_{\langle \xi_j(\omega) Ae_j \cup V \rangle_{j \neq i, j \in \sigma_1^c}} (Ae_i) \right\|^2 d\mu \\ &\leq \delta \int_{\Omega} \sum_{i \in \sigma_1^c} \left\| P_{\langle \xi_j(\omega) Ae_j \cup V \rangle_{j \in \sigma_1^c}} (Ae_i) \right\|^2 d\mu \end{aligned}$$

where the first equality used the fact that $\xi_i(\omega)$ and $\left\| P_{\langle \xi_j(\omega) Ae_j \cup V \rangle_{j \neq i, j \in \sigma_1^c}} (Ae_i) \right\|^2$ are independent (since the latter is a function of $(\xi_1, \dots, \xi_{i-1}, \xi_{i+1}, \dots, \xi_n)$ so that we may

split up the integral into a product) and the last line used the monotonicity of projection norms. By letting $W(\omega) = \{\xi_j(\omega)Ae_j : j \in \sigma_1^c\} \cup V$, we have

$$\begin{aligned}
 & \int_{\Omega} \sum_{i \in \sigma_1^c} \xi_i(\omega) \left\| P_{\langle \xi_j(\omega)Ae_j \cup V \rangle_{j \neq i, j \in \sigma_1^c}} (Ae_i) \right\|^2 d\mu \\
 & \leq \delta \int_{\Omega} \sum_{i \in \sigma_1^c} \left\| P_{\langle W(\omega) \rangle} (Ae_i) \right\|^2 d\mu \\
 & \leq \delta \int_{\Omega} \sum_{i=1}^n \left\| P_{\langle W(\omega) \rangle} (Ae_i) \right\|^2 d\mu \\
 & = \delta \int_{\Omega} \|P_{\langle W(\omega) \rangle} A\|_{\text{HS}}^2 d\mu \\
 & \leq \delta \|A\|^2 \int_{\Omega} \left(|\sigma_1| + \sum_{j \in \sigma_1^c} \xi_j(\omega) \right) d\mu \\
 & = \delta \|A\|^2 (|\sigma_1| + \delta |\sigma_1^c|)
 \end{aligned}$$

where we used the standard inequality $\|A\|_{\text{HS}}^2 \leq \|A\|^2 \text{rank}(A)$ and then made use of the fact that $\text{rank}(P_{\langle W(\omega) \rangle} A)$ is bounded above by the number of non-zero vectors in $\{\xi_i(\omega) : i \in \sigma_1^c\} \cup \{Ae_i : i \in \sigma_1\}$ which has the crude bound $|\sigma_1| + \sum_{j \in \sigma_1^c} \xi_j(\omega)$.

Since all functions involved are non-negative this yields:

$$\begin{aligned}
 & \int_{\Omega \setminus D} \sum_{i \in \sigma_1^c} \xi_i(\omega) \left\| P_{\langle \xi_j(\omega)Ae_j \cup V \rangle_{j \neq i, j \in \sigma_1^c}} (Ae_i) \right\|^2 d\mu \\
 & \leq \int_{\Omega} \sum_{i \in \sigma_1^c} \xi_i(\omega) \left\| P_{\langle \xi_j(\omega)Ae_j \cup V \rangle_{j \neq i, j \in \sigma_1^c}} (Ae_i) \right\|^2 d\mu \\
 & \leq \delta \|A\|^2 (|\sigma_1| + \delta |\sigma_1^c|)
 \end{aligned}$$

which, with (4), implies that there exists a point $\omega_0 \in \Omega \setminus D$ such that

$$\begin{aligned}
 \sum_{i \in \sigma(\omega_0)} \left\| P_{\langle Ae_j \cup V \rangle_{j \in \sigma(\omega_0) \setminus \{i\}}} (Ae_i) \right\|^2 &= \sum_{i \in \sigma_1^c} \xi_i(\omega_0) \left\| P_{\langle \xi_j(\omega_0)Ae_j \cup V \rangle_{j \neq i, j \in \sigma_1^c}} (Ae_i) \right\|^2 \\
 &\leq \frac{\delta \|A\|^2 (|\sigma_1| + \delta |\sigma_1^c|)}{1 - \mu(D)} \\
 &\leq \frac{\delta \|A\|^2 (|\sigma_1| + \delta |\sigma_1^c|)}{1 - 4(\delta |\sigma_1^c|)^{-1}}.
 \end{aligned}$$

The denominator can be dealt with by working with two separate cases:

Case 1: $\delta n < 32$. In this case since $d_1 = 1/(8 \cdot 32)$ we have that

$$d_1 \frac{n}{\|A\|^2} = \frac{n\delta}{32} < 1$$

so that $|\sigma_1| = 0$, and for any $\sigma_2 \subset \{1, \dots, n\}$, with $|\sigma_2| = 1$ the statement is vacuously true.

Case 2: $\delta n \geq 32$. Note that the assumption made on the columns of A imply that $\|A\|^2 \geq 1$. So since $d_1 < 1/2$ and $|\sigma_1| \leq d_1 n \|A\|^{-2}$ we must have that $|\sigma_1^c| \geq n/2$, i.e., $4(\delta|\sigma_1^c|)^{-1} \leq 8(\delta n)^{-1} \leq 1/3$. Therefore,

$$\sum_{i \in \sigma(\omega_0)} \left\| P_{\langle Ae_j \cup V \rangle_{j \in \sigma(\omega_0) \setminus \{i\}}} (Ae_i) \right\|^2 \leq \frac{\delta \|A\|^2 (|\sigma_1| + \delta |\sigma_1^c|)}{1 - 1/3} \leq \frac{3}{2} \delta \|A\|^2 (|\sigma_1| + \delta |\sigma_1^c|).$$

By definition of D , we also have:

$$|\sigma(\omega_0)| = \sum_{i \in \sigma_1^c} \xi_i(\omega_0) \geq \delta |\sigma_1^c|/2.$$

Let

$$\sigma_2 = \left\{ i \in \sigma(\omega_0) : \left\| P_{\langle Ae_j \cup V \rangle_{j \in \sigma(\omega_0) \setminus \{i\}}} (Ae_i) \right\| < 2 \|A\| \sqrt{\delta} \right\}.$$

Note that, given our choice of δ , σ_2 satisfies the desired conclusion. We just need to provide a lower bound on its size. Now,

$$\begin{aligned} \sum_{i \in \sigma(\omega_0) \setminus \sigma_2} \left\| P_{\langle Ae_j \cup V \rangle_{j \in \sigma(\omega_0) \setminus \{i\}}} (Ae_i) \right\|^2 &\leq \sum_{i \in \sigma(\omega_0)} \left\| P_{\langle Ae_j \cup V \rangle_{j \in \sigma(\omega_0) \setminus \{i\}}} (Ae_i) \right\|^2 \\ &\leq \frac{3}{2} \delta \|A\|^2 (|\sigma_1| + \delta |\sigma_1^c|). \end{aligned}$$

so, using the definition of σ_2 ,

$$4 \|A\|^2 \delta |\sigma(\omega_0) \setminus \sigma_2| \leq \frac{3}{2} \delta \|A\|^2 (|\sigma_1| + \delta |\sigma_1^c|).$$

Since $|\sigma_1| \leq d_1 n \|A\|^{-2}$ and $|\sigma(\omega_0)| \geq \delta |\sigma_1^c|/2$, we have

$$4 \|A\|^2 \delta |\sigma(\omega_0) \setminus \sigma_2| \leq \frac{3}{2} \delta \|A\|^2 \left(d_1 \frac{n}{\|A\|^2} + 2 |\sigma(\omega_0)| \right).$$

Solving this inequality for $|\sigma_2|$ yields

$$\begin{aligned} |\sigma_2| &\geq \frac{1}{4} |\sigma(\omega_0)| - \frac{3}{8} d_1 \frac{n}{\|A\|^2} \geq \delta \frac{|\sigma_1^c|}{8} - d_1 \frac{n}{\|A\|^2} = \frac{|\sigma_1^c|}{64 \|A\|^2} - \frac{n}{200 \|A\|^2} \\ &\geq \frac{n}{128 \|A\|^2} - \frac{n}{200 \|A\|^2} \geq \frac{n}{320 \|A\|^2} = d_1 \frac{n}{\|A\|^2}. \quad \square \end{aligned}$$

LEMMA 4.4. *For every $\sigma_1 \subset \{1, \dots, n\}$ with $|\sigma_1| \leq d_1 n \|A\|^{-2}$ there exists $\sigma_2 \subset \{1, \dots, n\} \setminus \sigma_1$ with $|\sigma_2| \geq (d_1/2) n \|A\|^{-2}$ such that for every $\{a_j\}_{j \in \sigma_1 \cup \sigma_2}$:*

$$\left\| \sum_{j \in \sigma_1 \cup \sigma_2} a_j Ae_j \right\| \geq \frac{1}{4\sqrt{2|\sigma_2|}} \sum_{j \in \sigma_2} |a_j|.$$

Proof. From Lemma 4.3 there exists $\sigma \subset \{1, \dots, n\} \setminus \sigma_1$ with $|\sigma| \geq d_1 n / \|A\|^2$, such that if for every $i \in \sigma$ we let $Ae_i = x_i$, then

$$\|P_{\langle Ae_k: k \in (\sigma_1 \cup \sigma) \setminus \{i\} \rangle} x_i\| < \frac{1}{\sqrt{2}}.$$

For every $i \in \sigma$, let $u'_i = x_i - P_{\langle Ae_k: k \in (\sigma_1 \cup \sigma) \setminus \{i\} \rangle} x_i$. Then, by orthogonality,

$$\|u'_i\|^2 = \|x_i\|^2 - \|P_{\langle Ae_k: k \in (\sigma_1 \cup \sigma) \setminus \{i\} \rangle} x_i\|^2 > 1 - \frac{1}{2} = \frac{1}{2}$$

and $\|u'_i\| \leq 1$. In addition, for $i \in \sigma_1 \cup \sigma$ and $j \in \sigma$ with $i \neq j$,

$$\begin{aligned} \langle x_i, u'_j \rangle &= \langle x_i, x_j - P_{\langle Ae_k: k \in (\sigma_1 \cup \sigma) \setminus \{j\} \rangle} x_j \rangle \\ &= \langle x_i, x_j \rangle - \langle x_i, P_{\langle Ae_k: k \in (\sigma_1 \cup \sigma) \setminus \{j\} \rangle} x_j \rangle \\ &= \langle x_i, x_j \rangle - \langle P_{\langle Ae_k: k \in (\sigma_1 \cup \sigma) \setminus \{j\} \rangle} x_i, x_j \rangle = 0. \end{aligned}$$

Similarly, for $i \in \sigma$

$$\begin{aligned} \langle x_i, u'_i \rangle &= \|x_i\|^2 - \langle x_i, P_{\langle Ae_k: k \in (\sigma_1 \cup \sigma) \setminus \{i\} \rangle} x_i \rangle \\ &= 1 - \|P_{\langle Ae_k: k \in (\sigma_1 \cup \sigma) \setminus \{i\} \rangle} x_i\|^2 > 1/2. \end{aligned}$$

For $i \in \sigma$ let $u_i = \|u'_i\|^{-1} u'_i$, so that $1 \geq \langle x_i, u_i \rangle > 1/2$.

If $E(X)$ denotes the expected value of the random variable X over $\{-1, 1\}^\sigma$ with the uniform probability measure, a simple calculation yields

$$\mathbb{E}\left(\left\|\sum_{i \in \sigma} \varepsilon_i u_i\right\|^2\right) = \sum_{i \in \sigma} \|u_i\|^2 = |\sigma|.$$

Thus if

$$\mathcal{E} = \left\{(\varepsilon_i)_{i \in \sigma} \in \{-1, 1\}^\sigma : \left\|\sum_{i \in \sigma} \varepsilon_i u_i\right\| \leq 2\sqrt{|\sigma|}\right\},$$

it follows then by Markov's inequality that

$$|\mathcal{E}| \geq \frac{3}{4} 2^{|\sigma|}.$$

By a theorem of Sauer and Shelah (see, e.g., [5, Page 144] or [19]), whenever k satisfies

$$|\mathcal{E}| > \sum_{i=0}^{k-1} \binom{|\sigma|}{i}, \tag{5}$$

then there exists a subset $\sigma_2 \subset \sigma$ of cardinality k such that for each tuple $(\varepsilon_i)_{i \in \sigma_2}$ there exists an extension $(\varepsilon_i)_{i \in \sigma}$ which belongs to \mathcal{E} . Note that (5) holds for $k \geq \frac{|\sigma|}{2}$ and therefore we may choose σ_2 with $|\sigma_2| \geq (d_1 n) / (2\|A\|^2)$ and $|\sigma_2| \geq |\sigma|/2$.

To see that σ_2 satisfies the conclusion, let $\{a_j\}_{j \in \sigma_1 \cup \sigma_2}$ be given. Define $\theta_i \in \{-1, 1\}$ for each $i \in \sigma_2$ so that $\theta_i a_i = |a_i|$. Then let $(\varepsilon_i)_{i \in \sigma}$ be an extension of $(\theta_i)_{i \in \sigma_2}$ that belongs to \mathcal{E} . It follows that

$$\begin{aligned} \left| \left\langle \sum_{j \in \sigma} \varepsilon_j u_j, \sum_{j \in \sigma_1 \cup \sigma_2} a_j x_j \right\rangle \right| &\leq \left\| \sum_{j \in \sigma} \varepsilon_j u_j \right\| \left\| \sum_{j \in \sigma_1 \cup \sigma_2} a_j x_j \right\| \leq 2\sqrt{|\sigma|} \left\| \sum_{j \in \sigma_1 \cup \sigma_2} a_j x_j \right\| \\ &\leq 2\sqrt{2|\sigma_2|} \left\| \sum_{j \in \sigma_1 \cup \sigma_2} a_j x_j \right\|. \end{aligned}$$

Hence

$$\begin{aligned} 2\sqrt{2}\sqrt{|\sigma_2|} \left\| \sum_{j \in \sigma_1 \cup \sigma_2} a_j x_j \right\| &\geq \left| \left\langle \sum_{j \in \sigma} \varepsilon_j u_j, \sum_{j \in \sigma_1 \cup \sigma_2} a_j x_j \right\rangle \right| \\ &= \sum_{j \in \sigma_2} |a_j| \langle x_j, u_j \rangle \geq \frac{1}{2} \sum_{j \in \sigma_2} |a_j|. \quad \square \end{aligned}$$

In the next step, we will require Khintchine's inequality which states the following. Let $\{1, -1\}^n$ be endowed with the uniform probability measure. For $\{b_\ell\}_{\ell=1}^n$ in \mathbb{R} we have

$$\mathbb{E} \left(\left| \sum_{\ell=1}^n \varepsilon_\ell b_\ell \right| \right) \geq \frac{1}{\sqrt{2}} \left(\sum_{\ell=1}^n |b_\ell|^2 \right)^{1/2}.$$

LEMMA 4.5. *For every $\sigma_1 \subset \{1, \dots, n\}$ with $|\sigma_1| \leq d_1 n \|A\|^{-2}$, there exists $\sigma_2 \subset \{1, \dots, n\} \setminus \sigma_1$ with $|\sigma_2| \geq (d_1/4)n \|A\|^{-2}$ such that for every $\{a_j\}_{j \in \sigma_1 \cup \sigma_2}$ in \mathbb{R} :*

$$\left\| \sum_{j \in \sigma_1 \cup \sigma_2} a_j A e_j \right\| \geq \frac{1}{16\sqrt{2}} \left(\sum_{j \in \sigma_2} |a_j|^2 \right)^{1/2}.$$

Proof. Consider the set σ supplied by applying Lemma 4.4 and denote $c' = 1/(4\sqrt{2})$. We need to establish a subset $\sigma_2 \subset \sigma$ of cardinality $|\sigma_2| \geq |\sigma|/2$ such that for any choice of coefficients in $\{a_j\}_{j \in \sigma_1 \cup \sigma_2}$:

$$\left\| \sum_{j \in \sigma_1 \cup \sigma_2} a_j A e_j \right\| \geq \frac{c'}{4} \left(\sum_{j \in \sigma_2} |a_j|^2 \right)^{1/2}.$$

Suppose, for contradiction, that such a subset does not exist. Let $A e_j = x_j$. Put $v_1 = \sigma$. Then there exists a vector $y_1 = \sum_{k \in v_1 \cup \sigma_1} b_{1,k} x_k$ such that $\|y_1\| < c'/4$, but $\sum_{k \in v_1} |b_{1,k}|^2 = 1$.

Assume that we have already constructed subsets $v_1 \supset v_2 \supset \dots \supset v_p$ with $|v_p| \geq |\sigma|/2$, and vectors $\{y_\ell\}_{\ell=1}^p$ such that $y_\ell = \sum_{k \in v_\ell \cup \sigma_1} b_{\ell,k} x_k$ and $\|y_\ell\| < c'/4$ and $\sum_{k \in v_\ell} |b_{\ell,k}|^2 = 1$, for $1 \leq \ell \leq p$. Consider the set

$$v_{p+1} = \left\{ k \in v_p : \sum_{\ell=1}^p |b_{\ell,k}|^2 < 1 \right\}.$$

If $|v_{p+1}| < |\sigma|/2$, then stop the procedure. On the other hand, if $|v_{p+1}| \geq |\sigma|/2$, then there exists a vector

$$y_{p+1} = \sum_{k \in v_{p+1} \cup \sigma_1} b_{p+1,k} x_k$$

such that $\|y_{p+1}\| < c'/4$ and $\sum_{k \in v_{p+1}} |b_{p+1,k}|^2 = 1$.

By the pigeon-hole principle, this algorithm must eventually terminate, say after m steps. Then

$$|v_{m+1}| < \frac{|\sigma|}{2}$$

and thus, for $k \in \sigma \setminus v_{p+1}$, we have that

$$\sum_{\ell=1}^m |b_{\ell,k}|^2 \geq 1$$

with the convention that $b_{\ell,k} = 0$ for those ℓ and k for which $b_{\ell,k}$ has not been defined.

Hence

$$m = \sum_{\ell=1}^m \sum_{k \in v_\ell} |b_{\ell,k}|^2 = \sum_{k \in \sigma} \sum_{\ell=1}^m |b_{\ell,k}|^2 \geq \sum_{k \in \sigma \setminus v_{m+1}} \sum_{\ell=1}^m |b_{\ell,k}|^2 \geq |\sigma \setminus v_{m+1}|$$

and this implies that $m \geq |\sigma|/2$.

On the other hand, we have

$$\begin{aligned} \frac{c' \sqrt{m}}{4} &> \left(\sum_{\ell=1}^m \|y_\ell\|^2 \right)^{\frac{1}{2}} = \left(\int \left\| \sum_{\ell=1}^m \varepsilon_\ell y_\ell \right\|^2 d\varepsilon \right)^{\frac{1}{2}} \geq \int \left\| \sum_{\ell=1}^m \varepsilon_\ell y_\ell \right\| d\varepsilon \\ &= \int \left\| \sum_{k \in \sigma \cup \sigma_1} \left(\sum_{\ell=1}^m \varepsilon_\ell b_{\ell,k} \right) x_k \right\| d\varepsilon \\ &\geq \frac{c'}{\sqrt{|\sigma|}} \sum_{k \in \sigma} \int \left| \sum_{\ell=1}^m \varepsilon_\ell b_{\ell,k} \right| d\varepsilon \end{aligned} \quad (6)$$

$$\geq \frac{c'}{\sqrt{2|\sigma|}} \sum_{k \in \sigma} \left(\sum_{\ell=1}^m |b_{\ell,k}|^2 \right)^{\frac{1}{2}} \quad (7)$$

where (6) follows from Lemma 4.4, and (7) follows from Khintchine's inequality.

However, the inductive construction implies

$$\sum_{\ell=1}^m |b_{\ell,k}|^2 \leq 2 \quad (8)$$

for all $k \in \sigma$. This is clear if $k \in v_{m+1}$ because of how we construct v_{m+1} , while if $k \in (v_p \setminus v_{p+1})$ for some $1 \leq p \leq m$, we have that

$$\sum_{\ell=1}^m |b_{\ell,k}|^2 = \sum_{\ell=1}^{p-1} |b_{\ell,k}|^2 + |b_{p,k}|^2 < 2.$$

It follows from (7) and (8) that

$$\begin{aligned} \frac{\sqrt{m|\sigma|}}{2} &> \sqrt{2} \sum_{k \in \sigma} \left(\sum_{\ell=1}^m |b_{\ell,k}|^2 \right)^{\frac{1}{2}} \geq \sum_{k \in \sigma} \left(\sum_{\ell=1}^m |b_{\ell,k}|^2 \right)^{\frac{1}{2}} \left(\sum_{\ell=1}^m |b_{\ell,k}|^2 \right)^{\frac{1}{2}} \\ &= \sum_{k \in \sigma} \sum_{\ell=1}^m |b_{\ell,k}|^2 = m. \end{aligned}$$

Thus $m < |\sigma|/4$ which contradicts the fact that $m \geq |\sigma|/2$. \square

We now prove Theorem 4.1.

Proof of Theorem 4.1. First apply Lemma 4.5 to obtain σ_2 that satisfies the desired conclusion for $d_2 = d_1/4$. We will next assume $|\sigma_1| \geq d_2 n \|A\|^{-2}$ and $d_1 d_2 n \|A\|^{-4} \geq 1$.

Let $d_3 = d_2^2$. Choose $\tilde{\sigma}_2 \subset \sigma_2$ with the largest possible cardinality that satisfies the implicit bound $|\tilde{\sigma}_2| \leq d_1(|\sigma_1| + |\tilde{\sigma}_2|)/\|A\|^2$ (such sets exist; the empty set is one such example). Define $n' = |\sigma_1| + |\tilde{\sigma}_2|$ and note that

$$n' \geq |\sigma_1| \geq d_2 \frac{n}{\|A\|^2}.$$

Since $|\sigma_2| \geq d_2 n / \|A\|^2$, the set σ_2 is sufficiently large to have allowed us to choose $\tilde{\sigma}_2$ such that

$$\begin{aligned} |\tilde{\sigma}_2| &\geq \lfloor d_1 d_2 n / \|A\|^4 \rfloor \geq (d_1 d_2 n / \|A\|^4) / 2 \quad (\text{because } d_1 d_2 n / \|A\|^4 \geq 1) \\ &\geq d_3 n / \|A\|^4. \end{aligned}$$

Now we define $V = \langle e_j : j \in \sigma_1 \cup \tilde{\sigma}_2 \rangle$ and $A' = A|_V$. An application of Lemma 4.5 to the matrix A' and the set $\tilde{\sigma}_2$, yields a subset $\tau_1 \subset \sigma_1$ such that

$$|\tau_1| \geq d_2 n' / \|A'\|^2 \geq d_2^2 n / \|A\|^4 \geq d_3 n / \|A\|^4,$$

and for any choice of scalars $\{a_j\}_{j \in \tau_1 \cup \tilde{\sigma}_2}$, we have

$$\left\| \sum_{j \in \tau_1 \cup \tilde{\sigma}_2} a_j A e_j \right\| \geq \frac{1}{16\sqrt{2}} \left(\sum_{j \in \tau_1} |a_j|^2 \right)^{1/2}. \quad (9)$$

As σ_2 was provided by Lemma 4.5 and $\tau_1 \subset \sigma_1$, $\tilde{\sigma}_2 \subset \sigma_2$ we also have

$$\left\| \sum_{j \in \tau_1 \cup \tilde{\sigma}_2} a_j A e_j \right\| \geq \frac{1}{16\sqrt{2}} \left(\sum_{j \in \tilde{\sigma}_2} |a_j|^2 \right)^{1/2}. \quad (10)$$

Recall that for any two non-negative numbers x, y we have $\max\{x, y\} \geq (1/\sqrt{2})(x^2 + y^2)^{1/2}$, which in conjunction with (9) and (10) yields the desired inequality. \square

The above theorem guarantees the existence of subsets of columns that can be “stitched” together without violating the minimal stretch property.

4.2. The proof of Theorem 1.4

Recall that an m -dimensional subspace U of \mathbb{R}^n is a λ -quadratic convex combination of U_{σ_1} and U_{σ_2} , where $\lambda \in [0, 1]$ and $\sigma_1 = \{i_1 < \dots < i_m\}$, $\sigma_2 = \{j_1 < \dots < j_m\}$ are disjoint subsets of $\{1, \dots, n\}$, such that U is spanned by the orthonormal sequence $u_k = \lambda^{1/2} e_{i_k} + (1 - \lambda)^{1/2} e_{j_k}$, $k = 1, \dots, m$.

In order to prove Theorem 1.4, we will combine Theorem 4.1 with a continuous traversal between disjoint subsets of the basis via quadratic convex combinations.

LEMMA 4.6. *Let $A : [a, b] \rightarrow M_{n \times n}$ be a continuous matrix function and let $\sigma_1, \sigma_2 \subset \{1, \dots, n\}$ such that $|\sigma_1| = |\sigma_2|$ and $\sigma_1 \cap \sigma_2 = \emptyset$. Let $U = \langle e_i : i \in \sigma_1 \cup \sigma_2 \rangle$ and suppose there exists a $c > 0$ such that $m_{A(t)|_U} \geq c$ for all $t \in [a, b]$. Let $\{U(t)\}_{t \in [a, b]}$ be such that, for each $t \in [a, b]$, $U(t)$ is the $(1 - (t - a)/(b - a))$ -quadratic convex combination of U_{σ_1} and U_{σ_2} . Then $\{U(t)\}_{t \in [a, b]}$ is a continuous choice of subspaces of \mathbb{R}^n such that $U(a) = U_{\sigma_1}$, $U(b) = U_{\sigma_2}$, and $m_{A(t)|_{U(t)}} \geq c$ for all $t \in [a, b]$.*

Proof. Write $\sigma_1 = \{i_1 < \dots < i_m\}$, $\sigma_2 = \{j_1 < \dots < j_m\}$ so that for every $t \in [a, b]$,

$$u_k(t) = \left(1 - \frac{t-a}{b-a}\right)^{1/2} e_{i_k} + \left(\frac{t-a}{b-a}\right)^{1/2} e_{j_k}, \quad k = 1, \dots, m$$

is an orthonormal basis for $U(t)$. As this is a continuous choice of orthonormal basis, by Lemma 2.4, $\{U(t)\}_{t \in [a, b]}$ is a continuous choice of subspaces and clearly $U(a) = U_{\sigma_1}$, $U(b) = U_{\sigma_2}$.

To complete the proof, for $t \in [a, b]$ note that

$$\begin{aligned} m_{A(t)|_{U(t)}} &= \inf\{\|A(t)x\| : x \in U(t), \|x\| = 1\} \\ &\geq \inf\{\|A(t)x\| : x \in U, \|x\| = 1\} \quad (\text{as } U(t) \subset U) \\ &= m_{A(t)|_U} \geq c. \quad \square \end{aligned}$$

We now have all the tools at our disposal to prove Theorem 1.4, which is restated for convenience.

THEOREM 4.7. *There exist universal constants $c = 1/33$ and $d = 2^{-21}$ such that for all continuous matrix functions $A : \mathbb{R} \rightarrow M_{n \times n}$ with the property that $\|A(t)e_i\| = 1$ for all $t \in \mathbb{R}$ and $1 \leq i \leq n$, there exists a continuous family of m -dimensional subspaces $\{U(t)\}_{t \in \mathbb{R}}$ of \mathbb{R}^n with $m \geq dn/\Lambda^4$ where $\Lambda = \sup_{t \in \mathbb{R}} \|A(t)\|$ such that $\|A(t)v\| \geq c\|v\|$ for every $t \in \mathbb{R}$ and every $v \in U(t)$. Furthermore, each subspace $U(t)$ is a quadratic convex combination of disjoint basis vectors.*

Proof. Note that $c = 1/33$ is a perturbation of $1/32$ and $d = d_3$, which are from the conclusion of Theorem 4.1 and Remark 4.2. If $d_1 d_2 n / \Lambda^4 \leq 1$, then we may simply select $U(t) = \langle e_1 \rangle$, for all $t \in \mathbb{R}$. Hence, we may assume $d_1 d_2 n / \Lambda^4 \geq 1$. Note that

$$\left\lceil d_2 \frac{n}{\Lambda^2} \right\rceil = \left\lceil \frac{1}{4} d_1 \frac{n}{\Lambda^2} \right\rceil \leq d_1 \frac{n}{\Lambda^2},$$

since $d_1 n / \Lambda^2 \geq d_1 d_2 n / \Lambda^4 \geq 1$. Furthermore,

$$\left\lceil d_3 \frac{n}{\Lambda^4} \right\rceil \leq d_3 \frac{n}{\Lambda^4} + 1 \leq \frac{d_2}{2\Lambda^2} \left(d_2 \frac{n}{\Lambda^2} \right) + \frac{d_1}{\Lambda^2} \left(d_2 \frac{n}{\Lambda^2} \right) \leq 2d_1 \left(d_2 \frac{n}{\Lambda^2} \right) < \frac{1}{3} \left\lceil d_2 \frac{n}{\Lambda^2} \right\rceil. \quad (11)$$

By Lemma 3.4 we can find an increasing sequence of points $(t_i)_{i \in \mathbb{Z}}$ such that for all $i \in \mathbb{Z}$ and t in $[t_{i-1}, t_{i+1}]$, $\|A(t_i) - A(t)\| \leq \varepsilon = 1/32 - 1/33$. Moreover, by Theorem 4.1 applied to the matrix $A(t_0)$ and the empty set, there exists a subset $\sigma_0 \subset \{1, \dots, n\}$ with $|\sigma_0| = \lceil d_2 n / \Lambda^2 \rceil \leq d_1 n / \Lambda^2$ such that for any choice of coefficients $\{a_j\}_{j \in \sigma_0}$ we have

$$\left\| \sum_{j \in \sigma_0} a_j A(t_0) e_j \right\| \geq \frac{1}{16\sqrt{2}} \left(\sum_{j \in \sigma_0} |a_j|^2 \right)^{1/2} \geq \frac{1}{32} \left(\sum_{j \in \sigma_0} |a_j|^2 \right)^{1/2}.$$

By using $\|A(t_0) - A(t)\| \leq \varepsilon = 1/32 - 1/33$ we obtain that if $U = \langle e_j : j \in \sigma_0 \rangle$, then $m_{A(t)|_U} \geq 1/33$ for all $t \in [t_{-1}, t_1]$ (see the last part of Lemma 3.4).

We now focus on the interval $[t_0, \infty)$ as a symmetric argument can be applied to $(-\infty, t_0]$. In reality, an extra stitching needs to be performed at t_0 to concatenate the two solutions. We omit this as it is essentially contained in the ensuing argument.

Iteratively applying Theorem 4.1, for $i \geq 0$, we can obtain the sets with following properties:

- (i) $\sigma_{i+1} \subset \{1, \dots, n\} \setminus \sigma_i$ with $|\sigma_{i+1}| = \lceil d_2 n / \Lambda^2 \rceil \leq d_1 n / \Lambda^2$ such that, for $U = \langle e_j : j \in \sigma_{i+1} \rangle$, we have $m_{A(t)|U} \geq 1/33$ for $t \in [t_i, t_{i+2}]$.
- (ii) $\tilde{\sigma}_{i+1} \subset \sigma_{i+1}$ and $\tilde{\tau}_i \subset \sigma_i$ with $|\tilde{\sigma}_{i+1}| = |\tilde{\tau}_i| = \lceil d_3 n / \Lambda^4 \rceil$ such that, for $U = \langle e_j : j \in \tilde{\tau}_i \cup \tilde{\sigma}_{i+1} \rangle$, we have $m_{A(t)|U} \geq 1/33$ for $t \in [t_i, t_{i+1}]$.

Indeed, suppose σ_i has been constructed with $|\sigma_i| = \lceil d_2 n / \Lambda^2 \rceil \leq d_1 n / \Lambda^2$ such that, for $U = \langle e_j : j \in \sigma_i \rangle$, we have $m_{A(t)|U} \geq 1/33$ for $t \in [t_i, t_{i+2}]$. To obtain the sets σ_{i+1} , $\tilde{\tau}_i$ and $\tilde{\sigma}_{i+1}$, we apply Theorem 4.1 to the matrix $A(t_{i+1})$ and the set σ_i . Because $|\sigma_i| \leq d_1 n / \Lambda^2$, we first get $\sigma_{i+1} \subset \{1, \dots, n\}$ with $|\sigma_{i+1}| \geq d_2 n / \Lambda^2$ (which, after truncating, we may assume satisfies $|\sigma_{i+1}| = \lceil d_2 n / \Lambda^2 \rceil$) such that, for any choice of coefficients $\{a_j\}_{j \in \sigma_i \cup \sigma_{i+1}}$,


$$\left\| \sum_{j \in \sigma_i \cup \sigma_{i+1}} a_j A(t_{i+1}) e_j \right\| \geq \frac{1}{32} \left(\sum_{j \in \sigma_{i+1}} |a_j|^2 \right)^{1/2}.$$

Using $\|A(t_{i+1}) - A(t)\| \leq \varepsilon = 1/32 - 1/33$ we obtain that, if $U = \langle e_j : j \in \sigma_{i+1} \rangle$, then $m_{A(t)|U} \geq 1/33$ for all $t \in [t_i, t_{i+2}]$. Since $|\sigma_i| = \lceil d_2 n / \Lambda^2 \rceil \geq d_2 n / \Lambda^2$, the second part of Theorem 4.1 yields $\tilde{\tau}_i \subset \sigma_i$ and $\tilde{\sigma}_{i+1} \subset \sigma_{i+1}$ with $|\tilde{\tau}_i|, |\tilde{\sigma}_{i+1}| \geq d_3 n / \Lambda^4$ (which after truncating we may assume $|\tilde{\tau}_i| = |\tilde{\sigma}_{i+1}| = \lceil d_3 n / \Lambda^4 \rceil$) such that for any choice of coefficients $\{a_j\}_{j \in \tilde{\tau}_i \cup \tilde{\sigma}_{i+1}}$ we have

$$\left\| \sum_{j \in \tilde{\tau}_i \cup \tilde{\sigma}_{i+1}} a_j A(t_{i+1}) e_j \right\| \geq \frac{1}{32} \left(\sum_{j \in \tilde{\tau}_i \cup \tilde{\sigma}_{i+1}} |a_j|^2 \right)^{1/2}.$$

By using $\|A(t_{i+1}) - A(t)\| \leq \varepsilon = 1/32 - 1/33$ we obtain that, if $U = \langle e_j : j \in \tilde{\tau}_i \cup \tilde{\sigma}_{i+1} \rangle$, then $m_{A(t)|U} \geq 1/33$ for all $t \in [t_i, t_{i+1}]$.

To construct our continuous collection of subspaces, consider the interval $[t_i, t_{i+1}]$. Using $\langle e_j : j \in \tilde{\tau}_i \rangle$ at t_i and $\langle e_j : j \in \tilde{\sigma}_{i+1} \rangle$ at t_{i+1} , we will invoke Lemma 4.6 to construct a continuous collection of subspaces $\{U(t)\}_{t \in [t_i, t_{i+1}]}$, that has dimension $\lceil d_3 n / \Lambda^4 \rceil$ for every $i \geq 0$. A diagram of the construction has been provided below for the reader's convenience:



As indicated on the diagram there is no reason to suppose $\tilde{\tau}_{i+1} = \tilde{\sigma}_{i+1}$, so we must devise a way to “stitch” together the collection of subspaces at each endpoint, in order to construct a continuous collection of subspaces $\{U(t)\}_{t \in \mathbb{R}}$. We choose an $\eta_i \in (0, \frac{t_{i+1}-t_i}{2})$ and allow for another instance of “stitching” in the interval $[t_{i+1} - \eta_i, t_{i+1}]$. This allows for us to continuously switch between $[t_i, t_{i+1}]$ and $[t_{i+1}, t_{i+2}]$.

It is necessary to observe here that both $\tilde{\tau}_{i+1}$ and $\tilde{\sigma}_{i+1}$ are subsets of σ_{i+1} . Thus the minimal stretch property holds for $U = \langle e_j : j \in \tilde{\tau}_{i+1} \cup \tilde{\sigma}_{i+1} \rangle$. We would like to switch from $\langle e_j : j \in \tilde{\sigma}_{i+1} \rangle$ at $t_{i+1} - \eta_i$ to $\langle e_j : j \in \tilde{\tau}_{i+1} \rangle$ at t_{i+1} . To do this, it suffices to choose a new subset $\xi_{i+1} \subset \sigma_{i+1} \setminus (\tilde{\tau}_{i+1} \cup \tilde{\sigma}_{i+1})$ with $|\xi_{i+1}| = |\tilde{\tau}_{i+1}| = |\tilde{\sigma}_{i+1}| = \lfloor d_3 n / \Lambda^4 \rfloor$. This is possible because

$$|\tilde{\tau}_{i+1} \cup \tilde{\sigma}_{i+1}| \leq 2 \left\lceil d_3 \frac{n}{\Lambda^4} \right\rceil < \frac{2}{3} \left\lceil d_2 \frac{n}{\Lambda^2} \right\rceil = \frac{2}{3} |\sigma_{i+1}| \quad (\text{by (11)}).$$

We can now switch between the following subspaces for each $[t_i, t_{i+1}]$:

At t_i : Begin with $U(t_i) = \langle e_j : j \in \tilde{\tau}_i \rangle$.

At $t_{i+1} - \eta_i$: Switch to $U(t_{i+1} - \eta_i) = \langle e_j : j \in \tilde{\sigma}_{i+1} \rangle$ using Lemma 4.6.

At $t_{i+1} - \eta_i/2$: Switch to $U(t_{i+1} - \eta_i/2) = \langle e_j : j \in \xi_i \rangle$ using Lemma 4.6.

At t_{i+1} : Switch to $U(t_{i+1}) = \langle e_j : j \in \tilde{\tau}_{i+1} \rangle$ using Lemma 4.6. \square

5. Future research directions

In Theorem 1.4 the continuous choice of subspaces preserves a strong type of coordinate structure. This comes at a steep price that needs to be paid in the dimensional estimate, which is of the order $n/\|A\|^4$, compared to the point-wise order $n/\|A\|^2$ provided by Theorem 1.1. The reason is the iterative application of the Bourgain-Tzafriri type Theorem 4.1. Therefore, there is potential room for improvement in answering Problem 2 in the introduction, e.g., by lowering the exponent 4 to which $\|A\|$ is raised, while preserving some of the coordinate structure of the subspaces.

An alternative direction is the achievement of kindred results to Theorem 1.4 with proxies of stable rank, e.g., the p -stable rank or entropic rank (see Remark 3.8) of an continuous $A : \mathbb{R} \rightarrow M_{n \times n}$, that does not necessarily have unit-length columns. It may be possible to obtain a continuous choice of subspaces which preserve some coordinate structure on which A is well invertible and with dimensional estimates similar to those found in [21], [20], [18], and [17].

It is also important noting that the topology of the domain \mathbb{R} of the continuous matrix function plays a big role on how the dimensional estimate $n/\|A\|^4$ is obtained in the proof of Theorem 1.4. Increasing the dimension of the domain might require further applications of a Bourgain-Tzafriri type theorem and thus further worsening the exponent of the estimate $n/\|A\|^4$ even further. It would be interesting to prove results similar to Theorems 1.2 and 1.4 for continuous matrix functions of a multi-dimensional domain.

Acknowledgements. The fourth and fifth authors are greatly indebted to Professors Alfonso Gracia-Saz (1976–2021) and Inder Bir Passi (1939–2021), without whose unfaltering support and encouragement the authors would never have considered continued studies in mathematics and to whom far more is owed than can be expressed. This paper is dedicated to their memories.

The authors would like to thank the anonymous referee whose invaluable comments led to the simplification of some arguments as well as to a more robust connection to the existing theory of restricted invertibility theorems.

REFERENCES

- [1] S. N. AFRIAT, *Orthogonal and oblique projectors and the characteristics of pairs of vector spaces*, Proc. Cambridge Philos. Soc. **53**: 800–816, 1957.
- [2] J. ANDERSON, *Extensions, restrictions, and representations of states on C^* -algebras*, Trans. Amer. Math. Soc. **249** (2): 303–329, 1979.
- [3] A. D. ANDREW, *Perturbations of Schauder bases in the spaces $C(K)$ and L^p , $p > 1$* , Studia Math. **65** (3): 287–298, 1979.
- [4] J. BOURGAIN AND S. J. SZAREK, *The Banach-Mazur distance to the cube and the Dvoretzky-Rogers factorization*, Israel J. Math. **62** (2): 169–180, 1988.
- [5] J. BOURGAIN AND L. TZAFRIRI, *Invertibility of “large” submatrices with applications to the geometry of Banach spaces and harmonic analysis*, Israel J. Math. **57** (2): 137–224, 1987.
- [6] J. BOURGAIN AND L. TZAFRIRI, *Restricted invertibility of matrices and applications*, in Analysis at Urbana, vol. II (Urbana, IL, 1986–1987), vol. 138 of London Math. Soc. Lecture Note Ser., pages 61–107, Cambridge Univ. Press, Cambridge, 1989.
- [7] J. BOURGAIN AND L. TZAFRIRI, *On a problem of Kadison and Singer*, J. Reine Angew. Math. **420**: 1–43, 1991.
- [8] P. G. CASAZZA AND J. C. TREMAIN, *Revisiting the Bourgain-Tzafriri restricted invertibility theorem*, Oper. Matrices **3** (1): 97–110, 2009.
- [9] Y. DAI, A. HORE, S. JIAO, T. LAN, AND P. MOTAKIS, *Continuous factorization of the identity matrix*, Involve **13** (1): 149–164, 2020.
- [10] R. V. KADISON AND I. M. SINGER, *Extensions of pure states*, Amer. J. Math. **81**: 383–400, 1959.
- [11] N. J. LAUSTSEN, R. LECHNER, AND P. F. X. MÜLLER, *Factorization of the identity through operators with large diagonal*, J. Funct. Anal. **275** (11): 3169–3207, 2018.
- [12] R. LECHNER, *Factorization in mixed norm Hardy and BMO spaces*, Studia Math. **242** (3): 231–265, 2018.
- [13] R. LECHNER, P. MOTAKIS, P. F. X. MÜLLER, AND TH. SCHLUMPRECHT, *Strategically reproducible bases and the factorization property*, Israel J. Math. **238** (1): 13–60, 2020.
- [14] R. LECHNER, P. MOTAKIS, P. F. X. MÜLLER, AND TH. SCHLUMPRECHT, *The factorisation property of $\ell^\infty(X_k)$* , Math. Proc. Cambridge Philos. Soc. **171** (2): 421–448, 2021.
- [15] S. J. LEON, *Linear algebra with applications*, Macmillan, Inc., New York; Collier-Macmillan Publishers, London, 1980.
- [16] A. W. MARCUS, D. A. SPIELMAN, AND N. SRIVASTAVA, *Interlacing families II: Mixed characteristic polynomials and the Kadison-Singer problem*, Ann. of Math. (2), **182** (1): 327–350, 2015.
- [17] ADAM W. MARCUS, DANIEL A. SPIELMAN, AND NIKHIL SRIVASTAVA, *Interlacing families III: Sharper restricted invertibility estimates*, Israel J. Math. **247** (2): 519–546, 2022.
- [18] A. NAOR AND P. YOUSSEF, *Restricted invertibility revisited*, in A journey through discrete mathematics, pages 657–691, Springer, Cham, 2017.
- [19] N. SAUER, *On the density of families of sets*, J. Combinatorial Theory Ser. A, **13**: 145–147, 1972.

- [20] D. A. SPIELMAN AND N. SRIVASTAVA, *An elementary proof of the restricted invertibility theorem*, Israel J. Math. **190**: 83–91, 2012.
- [21] R. VERSHYNIN, *John's decompositions: selecting a large part*, Israel J. Math. **122**: 253–277, 2001.

(Received January 12, 2022)

Adrian Fan
Department of Mathematics, Evans Hall, 3840
University of California
Berkeley, CA 94720-3840
e-mail: adrianfan@berkeley.edu

Jack Montemurro
Department of Mathematics
University of Toronto
Toronto, M5S 2E4, Canada
e-mail: jack.montemurro@mail.utoronto.ca

Pavlos Motakis
Department of Mathematics and Statistics
York University
4700 Keele Street, Toronto, Ontario, M3J 1P3, Canada
e-mail: pmotakis@yorku.ca

Naina Praveen
Department of Mathematics
Ashoka University, Rajiv Gandhi Education City
P.O. Rai, Sonapat, Haryana 131 029, India
e-mail: naina.praveen@alumni.ashoka.edu.in

Alyssa Rusonik
Department of Mathematics
University of Toronto
Toronto, M5S 2E4, Canada
e-mail: alyssa.rusonik@mail.utoronto.ca

Paul Skoufranis
Department of Mathematics and Statistics
York University
4700 Keele Street, Toronto, Ontario, M3J 1P3, Canada
e-mail: pskoufra@yorku.ca

Noam Tobin
Department of Mathematics and Statistics
York University
4700 Keele Street, Toronto, Ontario, M3J 1P3, Canada
e-mail: nsnow1@my.yorku.ca

